

Recombinant Horse IL18 Protein, His-tagged

Cat. No. IL18-1011H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Horse IL18 Protein (Tyr37-Asn193) with His tag was expressed in E. coli.
Species	Horse
Source	E.coli
ProteinLength	Tyr37-Asn193
Description	<p>Enables cytokine activity. Involved in lipid homeostasis; positive regulation of cell differentiation; and positive regulation of cold-induced thermogenesis. Acts upstream of or within several processes, including interleukin-18-mediated signaling pathway; positive regulation of NIK/NF-kappaB signaling; and positive regulation of interferon-gamma production. Located in extracellular space. Is expressed in several structures, including back skin; dorsal aorta; gut; lymphatic system; and reproductive system.</p> <p>Used to study atopic dermatitis. Human ortholog(s) of this gene implicated in several diseases, including Kawasaki disease; allergic rhinitis; autoimmune disease (multiple); biliary atresia; and hepatitis B. Orthologous to human IL18 (interleukin 18).</p>
Form	Freeze-dried powder
Molecular Mass	Predicted Molecular Mass: 21.7 kDa; Accurate Molecular Mass: 21 kDa
Purity	> 95%

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	Positive Control; Immunogen; SDS-PAGE; WB.
Stability	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 centigrade for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Storage	Avoid repeated freeze/thaw cycles. Store at 2-8 centigrade for one month. Aliquot and store at -80 centigrade for 12 months.
Storage Buffer	100mM NaHCO ₃ , 500mM NaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Reconstitution	Reconstitute in 100mM NaHCO ₃ , 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	IL18 interleukin 18 (interferon-gamma-inducing factor) [<i>Equus caballus</i>]
Official Symbol	IL18
Synonyms	IL18; interleukin 18 (interferon-gamma-inducing factor); interleukin-18; IL-1 gamma; interleukin-1 gamma; IFN-gamma-inducing factor; interferon gamma inducing factor; interferon gamma-inducing factor; interferon-gamma-inducing factor; IGIF; IL-18;
Gene ID	100034216
mRNA Refseq	NM_001082512
Protein Refseq	NP_001075981


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID

Q9XSQ7



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA