

Active Recombinant Human Interleukin 2

Cat. No. IL2-464H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant IL-2 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 134 amino acids and having a molecular mass of 15,517 Daltons.
Species	Human
Source	E.coli
ProteinLength	22-153 aa
Description	Interleukin-2, also know as T-cell Growth Factor, is an immunomodulatory cytokine produced by stimulated T-cells. Its activity is signaled via interaction with the IL-2R receptor.
Form	Human IL-2 is lyophilized at 1 mg/ml in 10 mM NaCitrate, pH 3.0.
Purity	Purity Greater than 90% as determined by: Analysis by RP-HPLC. Reducing and non-reducing SDS-PAGE.
Bio-activity	Recombinant Human IL-2 is fully bioactive when compared to standard. The ED50 as measured by the dose-dependent proliferation of CTLL-2 cells is 0.064 ng/ml.
Physical Appearance	Sterile Filtered white lyophilized (freeze-dried) powder.
Solubility	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg"s/ml, which can be further diluted into other aqueous solutions.

Purity Greater than 90% as determined by

Analysis by RP-HPLC. Reducing and non-reducing SDS-PAGE.

Protein Content determined by

UV spectroscopy at 280 nm. Analysis by RP-HPLC calibrated against a known standard. Quantitation on SDS-PAGE against a known standard.

Endotoxin Level

Endotoxin level is <0.01ng/ug or <0.1EU/ug.

Amino acid sequence

MAPTSSSTKK TQLQLEHLLL DLQMILNGIN NYKNPKLTRM LTFKFYMPKK
ATELKHLQCL EEELKPLEEV LNLAQSKNFH LRPRDLISNI NVIVLELKGS
ETTFMCEYAD ETATIVEFLN RWITFSQSII STLT

Storage

Lyophilized product is very stable at -20 degrees. Reconstituted material should be aliquoted and frozen at -20. It is recommended to add a carrier protein (0.1% HSA or BSA) for long term storage.

GENE INFORMATION

Gene Name

[IL2 interleukin 2 \[Homo sapiens \]](#)

Official Symbol

[IL2](#)

Synonyms

IL-2; TCGF; lymphokine; T-cell growth factor; Aldesleukin, Interleukin-2; interleukin-2 precursor; involved in regulation of T-cell clonal expansion; IL2; interleukin 2

Gene ID

[3558](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_000586
Protein Refseq	NP_000577
MIM	147680
UniProt ID	P60568
Chromosome Location	4q26-q27
Pathway	Cytokine-cytokine receptor interaction; Jak-STAT signaling pathway; T cell receptor signaling pathway; Type I diabetes mellitus
Function	interleukin-2 receptor binding; kinase activator activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA