

## Active Recombinant Human Interleukin 2, His-tagged

Cat. No. IL2-8861H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human Interleukin 2 was produced in the endosperm tissue of barley grain ( <i>Hordeum vulgare</i> ).
<b>Species</b>	Human
<b>Source</b>	<i>Hordeum Vulgare</i>
<b>Description</b>	Interleukin 2 (IL-2) is a potent immunoregulatory growth factor produced by T-cells following stimulation by mitogens or allogens. IL-2 promotes growth and differentiation of various cells of the immune system, such as activated T-cells and B-cells, NK cells, lymphokine activated killer cells, monocytes and macrophages. It is essential for proper immune response, it may also be an important factor in the natural suppression of autoimmunity. IL-2 may be of use as an anti-tumor agent in cancer therapy.
<b>Predicted N Terminal</b>	His
<b>Form</b>	Sterile filtered through a 0.2 µm filter. Lyophilized from PBS, pH 7.2.
<b>Bio-activity</b>	Bioactivity of recombinant human IL-2 is assayed by measuring its dose dependent effect on proliferation of CTLL-2 cells. The ED50 for this effect is typically < 0.25 ng/ml, corresponding to a specific activity > 4 x 10 <sup>6</sup> U/mg.
<b>Molecular Mass</b>	Recombinant human IL-2 contains 133 amino acids and a 16 a.a. histidine-based tag for a total length of 149 a.a. and has a predicted molecular mass of 17.5 kDa. The

Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

	recombinant protein migrates with an apparent molecular mass of 22 kDa in SDS-PAGE.
<b>Endotoxin</b>	Endotoxin level is less than 0.05EU/μg as measured by turbidimetric kinetic assay.
<b>Purity</b>	>95% by SDS-PAGE gel analysis.
<b>Storage</b>	The lyophilized protein is stable at room temperature for few weeks. Reconstituted protein should be used immediately or stored in working aliquots at -20°C. Avoid repeated freeze-thaw cycles.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration of no less than 100 μg/ml. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

## GENE INFORMATION

<b>Gene Name</b>	IL2 interleukin 2 [ Homo sapiens ]
<b>Official Symbol</b>	IL2
<b>Synonyms</b>	IL2; interleukin 2; interleukin-2; IL 2; T cell growth factor; TCGF; aldesleukin; involved in regulation of T-cell clonal expansion; IL-2; lymphokine;
<b>Gene ID</b>	3558
<b>mRNA Refseq</b>	NM_000586
<b>Protein Refseq</b>	NP_000577
<b>MIM</b>	147680

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>UniProt ID</b>	P60568
<b>Chromosome Location</b>	4q26-q27
<b>Pathway</b>	Allograft rejection, organism-specific biosystem; Allograft rejection, conserved biosystem; Autoimmune thyroid disease, organism-specific biosystem; Autoimmune thyroid disease, conserved biosystem; Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem; Calcium signaling in the CD4+ TCR pathway, organism-specific biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem;
<b>Function</b>	carbohydrate binding; cytokine activity; glycosphingolipid binding; growth factor activity; interleukin-2 receptor binding; interleukin-2 receptor binding; kappa-type opioid receptor binding; kinase activator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA