

Active Recombinant Human Interleukin 20

Cat. No. IL20-500H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL-20 produced in E.Coli is a homodimeric, non-glycosylated polypeptide chain containing 2 x 153 amino acids and having a total molecular mass of 35,212 Dalton.
Species	Human
Source	E.coli
Description	The newly discovered cytokine interleukin-20 (IL-20) is structurally related to IL-10, yet it appears to be an autocrine factor for keratinocytes that regulates their participation in inflammation.
Form	Recombinant hIL-20 was lyophilized from a concentrated (1mg/ml) solution containing 100mM sodium chloride and 10mM sodium citrate buffer pH-3.
Purity	Purity Greater than 98% as determined by: Analysis by RP-HPLC. Reducing and non-reducing SDS-PAGE.
Bio-activity	Human IL-20 is fully biologically active when compared to standard. The Biological Acitivity was determined by the ability to activate STAT following receptor ligand interaction.
Physical Appearance	Sterile Filtered white lyophilized (freeze-dried) powder.
Solubility	Centrifuge vial before opening. When reconstituting the product, gently pipet and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/ml, which can be further diluted into other aqueous solutions.

Purity Greater than 98% as determined by

Analysis by RP-HPLC. Reducing and non-reducing SDS-PAGE.

Protein Content determined by

UV spectroscopy at 280 nm. Analysis by RP-HPLC calibrated against a known standard. Quantitation on SDS-PAGE against a known standard.

Endotoxin Level

Endotoxin level as measured by LAL is <0.01ng/ug or <0.1EU/ug.

Amino acid sequence

MLKTLNLGSC VIATNLQEIR NGFSEIRGSV QAKDGNIDIR ILRRTESLQD
TKPANRCCLL RHLRLYLDR VFKNYQTPDH YTLRKISSLA NSFLTICKDL
RLCHAHMTCH CGEEAMKKYS QILSHFEKLE PQAAVVKALG ELDILLQWME ETE

Storage

Lyophilized product is very stable at -20 degrees. Reconstituted material should be aliquoted and frozen at -20. It is recommended to add a carrier protein (0.1% HSA or BSA) for long term storage.

Pathways

Cytokine-cytokine receptor interaction; Jak-STAT signaling pathway

GENE INFORMATION

Gene Name

IL20 interleukin 20 [Homo sapiens]

Official Symbol

IL20

Synonyms

IL20; interleukin 20; IL-20; IL10D; ZCYTO10; MGC96907; OTTHUMP00000034618;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

four alpha helix cytokine; OTTHUMP00000034617; Four alpha helix cytokine Zcyto10

Gene ID 50604**mRNA Refseq** NM_018724**Protein Refseq** NP_061194**MIM** 605619**UniProt ID** Q9NYY1**Chromosome
Location** 1q32**Function** cytokine activity; interleukin-20 receptor binding Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA