

Recombinant Human Interleukin 22

Cat. No. IL22-555H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	IL22 Human Recombinant produced in E. Coli is a single, non-glycosylated homodimeric polypeptide chain containing 2 x 146 amino acids and having a total MW 33.607kDalton.
Species	Human
Source	Human
Description	Interleukin-22 (IL22) is a protein that in humans is encoded by the IL22 gene. IL22 is a member of a group of cytokines called the IL-10 family or IL-10 superfamily (including IL-19, IL-20, IL-24, and IL-26), a class of potent mediators of cellular inflammatory responses. It shares use of IL-10R2 in cell signaling with other members of this family, IL-10, IL-26, IL-28A/B and IL-29.
Form	Lyophilized. Each mg contains 1.6 mg mannitol and 1.2 mg HSA.
Purity	Greater than 97.0% by SDS-PAGE gel and RF-HPLC.
Sequence	The sequence of the first five N-terminal amino acids is Met-Ala-Pro-Ile-Ser.
Molecular Weight	33.607kDalton
Biological Activity	Biological Activity was determined by the ability to activate STAT following receptor ligand interaction.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin Level	Less than 0.1 ng per g (1 EU/g) of protein.
Reconstitution	Centrifuge vial prior to opening. First add sterile water to the vial to fully solubilize the protein to a concentration not less than 100 g/ml. After complete solubilization of the protein, it can be further diluted to other aqueous solutions.
Storage	Lyophilized samples are stable at room temperature for 3 weeks, but it is recommended to store them desiccated at -20°C to -80°C. Upon reconstitution, protein should be stored at 2-4°C for one week and for six months at -20°C to -80°C. Add a carrier protein (0.1% HSA or BSA) as a stabilizer for long term storage. Avoid repeated freeze-thaw cycles.
Official Symbol	IL22

GENE INFORMATION

Gene Name	IL22interleukin 22 [Homo sapiens]
Synonyms	IL22; interleukin 22; TIFa; IL21; IL-21; IL-22; ILTIF; IL-TIF; IL-D110; zcyto18; ZCYTO183; MGC79382; MGC79384; TIFIL-23; interleukin 21; interleukin-22; cytokine Zcyto18; OTTHUMP00000240117; IL-10-related T-cell-derived inducible factor; IL-10-related T-cell-derived-inducible factor
Gene ID	50616
mRNA Refseq	NM_020525
Protein Refseq	NP_065386
MIM	605330


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q9GZX6
Chromosome Location	12q15
Pathway	Cytokine-cytokinereceptor interaction; Jak-STAT signaling pathway
Function	cytokineactivity; interleukin-22 receptor binding

Crystallographic structure of IL-22 (rainbow colored, N-terminus blue, C-terminus red) complexed with the IL-22R1 (magenta).



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA