

# Active Recombinant Human IL2RA protein, Fc/Avi-tagged, Biotinylated

**Cat. No.** IL2RA-051H    **Lot. No.** (See product label)

## SPECIFICATION

<b>Product Overview</b>	Biotinylated Recombinant Human IL2RA(Glu22-Cys213) protein, fused to Fc/Avi tag at the C-terminus, was expressed in HEK293 cells .
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	Glu22-Cys213
<b>Description</b>	<p>IL-2 receptor alpha (IL2R alpha), also known as CD25, is a 55 kDa type I membrane glycoprotein that belongs to the family of cytokine receptors that utilize the common gamma chain subunit (gamma c). Human IL-2R alpha cDNA encodes a 213 amino acid (aa) precursor with a 21 aa signal peptide and a 192 aa extracellular region. The ECD of Human IL-2R alpha shares a 59% amino acid sequence identity with the ECD of mouse and rat IL2R alpha, respectively. IL-2R alpha is primarily expressed on activated T cells and on regulatory T cells (Treg) (1-3). IL-2R beta (CD122) and gamma c (IL-2R gamma /CD132) dimerize to form a constitutively expressed intermediate affinity IL-2 receptor (4, 5). By itself, IL-2R alpha binds IL-2 with low affinity. IL-2R alpha makes no contacts with IL-2R beta or gamma c, and only minor changes are observed in the IL-2 structure in response to receptor binding. These findings support the principal role of IL-2R alpha to deliver IL-2 to the signaling complex and act as regulator of signal transduction (6, 7). A soluble form of IL-2R alpha can be generated by proteolytic cleavage of the cell surface</p>

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

receptor, rendering the T cell unresponsive to IL-2 (8, 9). Increased serum levels of soluble IL-2R alpha are found in some cancers and immune disorders (10). IL-2R alpha is required for activation induced cell death (AICD) of naive T cells, a mechanism responsible for deleting autoreactive T cell clones (11, 12). IL-2R alpha is also required for the development of CD4+CD25+ Treg which suppresses autoreactive CD4+ T cells, thereby contributing to peripheral T cell homeostasis (11-13).

<b>Predicted N Terminal</b>	Glu22
<b>Form</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.
<b>Bio-activity</b>	Measured by its binding ability in a functional ELISA. When Recombinant Human IL2 is immobilized at 1 µg/mL (100 µ/well), Biotinylated Recombinant Human IL2RA Fc Chimera Avi-tag binds with an ED50 of 50-450 ng/mL.
<b>Molecular Mass</b>	69-77 kDa, under reducing conditions
<b>Endotoxin</b>	<0.10 EU per 1 µg of the protein by the LAL method.
<b>Purity</b>	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
<b>Applications</b>	Bioactivity
<b>Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Reconstitution** Reconstitute at 500 µg/mL in PBS.

**Conjugation** Biotin

## GENE INFORMATION

**Gene Name** [IL2RA interleukin 2 receptor, alpha \[ Homo sapiens \]](#)

**Official Symbol** [IL2RA](#)

**Synonyms** IL2RA; interleukin 2 receptor, alpha; IL2R; interleukin-2 receptor subunit alpha; CD25; p55; IL2-RA; IL-2-RA; TAC antigen; IL-2R subunit alpha; IL-2 receptor subunit alpha; TCGFR; IDDM10;

**Gene ID** [3559](#)

**mRNA Refseq** [NM\\_000417](#)

**Protein Refseq** [NP\\_000408](#)

**MIM** [147730](#)

**UniProt ID** [P01589](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

### SDS-PAGE



2 µg/lane Protein was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining.

### Binding Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA