

GMP Recombinant Human IL3 protein

Cat. No. IL3-132HG Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL3 protein was expressed in Escherichia coli. This product is produced under cGMP guidelines.
Species	Human
Source	E.coli
ProteinLength	133
Description	Interleukin-3 (IL-3) is an interleukin, a type of biological signal (cytokine) which is encoded by the IL-3 gene located on chromosome 5 and produced primarily by activated T cells beside human thymic epithelial cells, activated murine mast cells, murine keratinocytes and neurons/astrocytes. The protein acts in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. The human IL-3 reported to be a monomer, as it is known, contains 133 amino acids residues which is a single non-glycosylated polypeptide. Specifically, human and murine IL-3 share low homology and it does not show activity on murine cells.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4, with 0.02 % Tween-20.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using human TF-1 cells is less than 0.1 ng/ml, corresponding to a specific activity of > 1.0 ×

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	10 ⁷ IU/mg.
Molecular Mass	Approximately 15.0 kDa, a single non-glycosylated polypeptide chain containing 133 amino acids.
AA Sequence	APMTQTTSLKTSWVNCNMIIDEIITHLKQPPLPLLDNFNNLNGEDQQDILMENNLRPNL EAFNRAVKSLQNASAIESILKNLLPCLPLATAAPTRHPIHIKDGDWNEFRRKLTFLYKLT LENAQAQQTTLSLAIF
Endotoxin	Less than 0.01 EU/g of rHuIL-3 GMP as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	IL3
Official Symbol	IL3
Synonyms	IL3; interleukin 3 (colony-stimulating factor, multiple); interleukin-3; hematopoietic

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

growth factor; IL 3; mast cell growth factor; MCGF; MGC79398; MGC79399; MULTI-CSF; multilineage colony stimulating factor; P cell stimulating factor; mast-cell growth factor; P-cell stimulating factor; P-cell-stimulating factor; multilineage-colony-stimulating factor; multipotential colony-stimulating factor; IL-3; MULTI-CSF;

Gene ID [3562](#)


mRNA Refseq [NM_000588](#)

Protein Refseq [NP_000579](#)

MIM [147740](#)

UniProt ID [P08700](#)

SDS-PAGE of IL3-132HG




 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

HPLC of IL3-132HG



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA