

Active Recombinant Mouse IL3 protein, His-tagged

Cat. No. IL3-88M Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Mouse IL3 protein(P01586)(33~166 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Mouse
Source	HEK293
ProteinLength	33-166 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Interleukin 3 is an interleukin, a type of biological signal that can improve the body's natural response to disease as part of the immune system. It acts by binding to the interleukin 3 receptor IL-3 synergizes with other cytokines to stimulate the growth of immature progenitor cells of all lineages, and is therefore a multi-lineage colony-stimulating factor (CSF). It prevents cell death and promotes the survival of macrophages, mast cells, and megakaryocytes. To test the effect of IL-3 on cell proliferation, TF-1 cells were seeded into triplicate wells of 96-well plates at a density of 5,000 cells/well with 2% serum standard 1640 which contains various concentrations of recombinant mouse IL-3. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 µof CCK-8 solution was added to each well of the plate, then the absorbance at 450 nm was measured using a microplate reader after

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

incubating the plate for 2-4 hours at 37°C. Proliferation of TF-1 cells after incubation with IL-3 for 72h observed by inverted microscope was shown. Cell viability was assessed by CCK-8(Cell Counting Kit-8) assay after incubation with recombinant IL-3 for 72h. It was obvious that IL-3 significantly increased cell viability of TF-1 cells. The ED50 is 3.79 ng/ml.

Molecular Mass	16.7kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	IL3 interleukin 3 [<i>Mus musculus</i>]
Official Symbol	IL3
Synonyms	IL3; interleukin 3; interleukin-3; mast cell growth factor; P-cell-stimulating factor; hematopoietic growth factor; multipotential colony-stimulating factor; BPA; PSF; HCGF; IL-3; MCGF; Csfmu;
Gene ID	16187
mRNA Refseq	NM_010556

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Protein Refseq

NP_034686

SDS-PAGE



Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA