

Recombinant Human IL33 protein

Cat. No. IL33-138H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL33 protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	159
Description	IL-33, encoded by IL-33 gene located on the Chr.9 in humans, is a 30 kDa proinflammatory protein belonging to the IL-1 superfamily and it shares less than 20 % a.a. sequence identity with other members. IL-33 secreted by high endothelial venules at high levels, which is found in tonsils, peyer patches and mesenteric lymph nodes, but not in placenta. It is upregulated in arterial smooth muscle, dermal fibroblasts, and keratinocytes following IL1 α or IL1 β stimulation. It elicits its biological effects by interacting with IL1RL1/ST2 and its stimulation recruits MYD88, IRAK1, IRAK4, and TRAF6, followed by phosphorylation of MAPK3/ERK1 and/or MAPK1/ERK2, MAPK14, and MAPK8. IL-33 mature protein has 52-58 % a.a. sequence identity with mouse and rat IL-33.
Form	Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM PB, 150 mM NaCl, 1mM EDTA, pH7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using murine D10S cells is less than 0.05 ng/ml, corresponding to a specific activity of > 2.0 \times

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	10 ⁷ IU/mg.
Molecular Mass	Approximately 17.9 kDa, a single non-glycosylated polypeptide chain containing 159 amino acids.
AA Sequence	SITGISPITEYLASLSTYNDQSITFALEDESYEIYVEDLKKDEKKDKVLLSYYESQHPNS ESGDGVDGKMLMVTLSPDKDFWLHANNKEHSVELHKCEKPLPDQAFFVLHNMHSN CVSFECKTDPGVFIGVKDNHLALIKVDSSENLCTENILFKLSET
Endotoxin	Less than 1 EU/g of rHuIL-33 as determined by LAL method.
Purity	>97% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	IL33
Official Symbol	IL33
Synonyms	IL33; interleukin 33; C9orf26, chromosome 9 open reading frame 26 (NF HEV);

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

interleukin-33; DKFZp586H0523; DVS27; DVS27 related protein; IL1F11; interleukin 1 family; member 11; NF HEV; nuclear factor for high endothelial venules; IL-33; IL-1F11; DVS27-related protein; interleukin-1 family member 11; nuclear factor from high endothelial venules; NF-HEV; NFEHEV; C9orf26; RP11-575C20.2;

Gene ID [90865](#)


mRNA Refseq [NM_001199640](#)

Protein Refseq [NP_001186569](#)

MIM [608678](#)

UniProt ID [O95760](#)

SDS-PAGE of IL33-138H



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA