

Recombinant Human IL4 Protein

Cat. No. IL4-514H Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant human IL4 protein

Species Human

ProteinLength 153

Description

The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. IL4 is considered an important cytokine for tissue repair, counterbalancing the effects of proinflammatory type 1 cytokines, however, it also promotes allergic airway inflammation. Moreover, IL-4, a type 2 cytokine, mediates and regulates a variety of human host responses such as allergic, anti-parasitic, wound healing, and acute inflammation. This cytokine has been reported to promote resolution of neutrophil-mediated acute lung injury. In an allergic response, IL-4 has an essential role in the production of allergen-specific immunoglobulin (Ig) E. This pro-inflammatory cytokine has been observed to be increased in COVID-19 (Coronavirus disease 2019) patients, but is not necessarily associated with severe COVID-19 pathology. Two alternatively spliced transcript

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

variants of this gene encoding distinct isoforms have been reported.

Form	Lyophilized
AA Sequence	MGLTSQLLPPLFFLLACAGNFVHGKCDITLQEIIKTLNSLTEQKTLCTELTVTDIFAAS KNTTEKETFCRAATVLRQFYSHHEKDTRCLGATAQQFHRHKQLIRFLKRLDRNLWG LAGLNSCPVKEANQSTLENFLERLKTIMREKYSKCSS
Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.

GENE INFORMATION

Gene Name	IL4 interleukin 4 [Homo sapiens (human)]
Official Symbol	IL4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Synonyms	IL4; interleukin 4; interleukin-4; B cell growth factor 1; B_cell stimulatory factor 1; BCGF 1; BCGF1; BSF1; IL 4; lymphocyte stimulatory factor 1; MGC79402; binetrakin; pitrakinra; IL-4; BSF-1; BCGF-1;
Gene ID	3565
mRNA Refseq	NM_000589
Protein Refseq	NP_000580
MIM	147780
UniProt ID	P05112

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA