

## Active Recombinant Human IL4,His-tagged

Cat. No. IL4-8863H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Human IL4 was produced in the endosperm tissue of barley grain ( <i>Hordeum vulgare</i> ).
<b>Species</b>	Human
<b>Source</b>	Hordeum Vulgare
<b>Description</b>	IL-4 (Interleukin 4, B cell stimulatory factor) is a pleiotropic cytokine produced by activated T cells, mast cells and basophils. IL-4 has many different functions including promoting proliferation and differentiation of immunologically competent cells. One of its main functions is regulating differentiation of naive CD4+ T cells into helper Th2 cells, the other main function is regulating B cells IgE and IgG1 production. Human IL-4 shows 50% homology with mouse IL-4, but its activities are species-specific and human IL-4 shows no activity on murine and rat cells.
<b>Predicted N Terminal</b>	His
<b>Form</b>	Sterile filtered through a 0.2 µm filter. Lyophilized from PBS, pH7,2
<b>Bio-activity</b>	The bioactivity of recombinant human IL-4 was determined by its dose-dependent effects on the proliferation of TF-1 cells. The ED50 value for this effect is typically = 0.2 ng/ml, corresponding to a specific activity of = 5 x 10e6 units/mg.
<b>Molecular Mass</b>	Recombinant human IL-4 contains 130 amino acids and a 16 a.a. Histidine-based tag for a total length of 146 a.a. and has a predicted molecular mass of 17.2 kDa. The

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	recombinant protein migrates as two bands with an apparent molecular mass of 23 kDa and 24 kDa in SDS-PAGE.
<b>Endotoxin</b>	Endotoxin level is less than 0.05EU/μg as measured by turbidimetric kinetic assay.
<b>Purity</b>	>95% by SDS-PAGE gel analysis.
<b>Storage</b>	The lyophilized protein is stable at room temperature for few weeks. Reconstituted protein should be used immediately or stored in working aliquots at -20°C. Avoid repeated freeze-thaw cycles.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration of no less than 100 μg/ml. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">IL4 interleukin 4 [ Homo sapiens ]</a>
<b>Official Symbol</b>	IL4
<b>Synonyms</b>	IL4; interleukin 4; interleukin-4; B cell growth factor 1; B_cell stimulatory factor 1; BCGF 1; BCGF1; BSF1; IL 4; lymphocyte stimulatory factor 1; MGC79402; binetrakin; pitrakinra; IL-4; BSF-1; BCGF-1;
<b>Gene ID</b>	<a href="#">3565</a>
<b>mRNA Refseq</b>	<a href="#">NM_000589</a>
<b>Protein Refseq</b>	<a href="#">NP_000580</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>MIM</b>	147780
<b>UniProt ID</b>	P05112
<b>Chromosome Location</b>	5q23-q31
<b>Pathway</b>	Allograft rejection, organism-specific biosystem; Allograft rejection, conserved biosystem; Asthma, organism-specific biosystem; Asthma, conserved biosystem; Autoimmune thyroid disease, organism-specific biosystem; Autoimmune thyroid disease, conserved biosystem; CD40/CD40L signaling, organism-specific biosystem;
<b>Function</b>	cytokine activity; growth factor activity; interleukin-4 receptor binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA