

Recombinant Human IL4 protein

Cat. No. IL4-99H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL4 protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	129
Description	Interleukin-4 (IL-4) is a pleiotropic cytokine that induces differentiation of naive helper T cells (Th0 cells) to Th2 cells and produced by by mast cells, activated T cells and bone marrow stromal cells. It has many biological roles, including the stimulation of activated B-cell and T-cell proliferation, and the differentiation of CD4+ T-cells into Th2 cells. In addition, It enhances both secretion and cell surface expression of IgE and IgG1 and also regulates the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. The human IL-4 has a compact, globular fold, stabilised by 3 disulphide bonds. The human IL-4 shares about 40% aa sequence identity with mouse/rat IL-4 and they are species-specific in their activities.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using human TF-1 cells is less than 0.2 ng/ml, corresponding to a specific activity of > 5.0 × 10 ⁶ IU/mg.
Molecular Mass	Approximately 15.0 kDa, a single non-glycosylated polypeptide chain containing 129

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	amino acids.
AA Sequence	HKCDITLQEIIKTLNSLTEQKTLCTELTVTDIFAASKNTTEKETFCRAATVLRQFYSHHE KDTRCLGATAQQFHRHKQLIRFLKRLDRNLWGLAGLNSCPVKEANQSTLENFLERL KTIMREKYSKCSS
Endotoxin	Less than 1.0 EU/g of rHuIL-4 as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.
GENE INFORMATION	
Gene Name	IL4
Official Symbol	IL4
Synonyms	IL4; interleukin 4; interleukin-4; B cell growth factor 1; B_cell stimulatory factor 1; BCGF 1; BCGF1; BSF1; IL 4; lymphocyte stimulatory factor 1; MGC79402; binetrakin; pitrakinra; IL-4; BSF-1; BCGF-1;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 3565


mRNA Refseq NM_000589

Protein Refseq NP_000580

MIM 147780

UniProt ID P05112

SDS-PAGE of IL4-99H



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA