

Recombinant Active Human IL7 Protein, His-tagged(C-ter)

Cat. No. IL7-214H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Active Human IL7 Protein with His tag (C-ter) was expressed in E. coli.
Species	Human
Source	E.coli
Description	<p>The protein encoded by this gene is a cytokine important for B and T cell development. This cytokine and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-stimulating factor. This cytokine is found to be a cofactor for V(D)J rearrangement of the T cell receptor beta (TCRB) during early T cell development. This cytokine can be produced locally by intestinal epithelial and epithelial goblet cells, and may serve as a regulatory factor for intestinal mucosal lymphocytes. Knockout studies in mice suggested that this cytokine plays an essential role in lymphoid cell survival. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional splice variants have been described but their presence in normal tissues has not been confirmed.[provided by RefSeq, Dec 2010]</p>
Form	Powder
Bio-activity	Measured in a cell proliferation assay using PHA-activated human peripheral blood lymphocytes (PBMC). The ED50 for this effect is < 0.8 ng/mL. The specific activity of recombinant human IL-7 is > 7 x 10 ⁸ IU/mg.
AA Sequence	MDCDIEGKDGKQYESVLMVSIQQLLDISMKEIGSNCLNNEFNFFKRHICDANKEGMFL

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

FRAARKLRQFLKMNSTGDFDLHLLKVSEGTILLNCTGQVKGRKPAALGEAQPTKSL
EENKSLKEQKKLNDLCFLKRLLEIKTCWNKILMGTKEH

Endotoxin	Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.
Purity	> 95% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.

GENE INFORMATION

Gene Name	IL7 interleukin 7 [Homo sapiens]
Official Symbol	IL7
Synonyms	IL7; interleukin 7; interleukin-7; IL 7; IL-7;
Gene ID	3574

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


mRNA Refseq [NM_000880](#)

Protein Refseq [NP_000871](#)

MIM [146660](#)

UniProt ID [P13232](#)

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA