

Active Recombinant Human INHBA, Animal Free

Cat. No. INHBA-131H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human Activin A is a 27.4 kDa disulphide-linked homodimers of two beta-A chains, each containing 116 amino residues. Human recombinant protein expressed in <i>Nicotiana benthamiana</i> . Recombinant human Activin A contains a 6-His-tag at the N-terminal end, is produced by transient expression in non-transgenic plants and is purified by sequential chromatography (FPLC). This product contains no animal-derived components or impurities. Animal free product.
Species	Human
Source	<i>Nicotiana Benthamiana</i>
ProteinLength	116 a.a.
Description	Activins are homodimers or heterodimers of the various beta subunit isoforms, belonging to the TGFbeta family. Mature Activin A has two 116 amino acids residues betaA subunits (betaA-betaA). Activin exhibits a wide range of biological activities, including mesoderm induction, neural cell differentiation, bone remodelling, haematopoiesis, and reproductive physiology. Activins plays a key role in the production and regulation of hormones such as FSH, LH, GnRH and ACTH. Cells known to express Activin A include fibroblasts, endothelial cells, hepatocytes, vascular smooth muscle cells, macrophages, keratinocytes, osteoclasts, bone marrow monocytes, prostatic epithelium, neurons, chondrocytes, osteoblasts, Leydig cells, Sertoli cells, and ovarian granulosa cells.As with other members of the super-family, Activins interact with two types of cell surface trans-membrane receptors (Types I and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	<p>II) which have intrinsic serine/threonine kinase activities in their cytoplasmic domains, Activin type 1 receptors, ACVR1, ACVR1B, ACVR1C and Activin type 2 receptors, ACVR2A, ACVR2B. The biological activity of Activin A can be neutralized by inhibins and by the diffusible TGF-B antagonist, Follistatin.</p>
Form	Lyophilized from a Tris HCl 0.05M buffer at pH 7.4.
Bio-activity	The biological activity of Activin A is measured by its ability to inhibit mouse plasmacytoma cell line (MPC-11) cells proliferation ([³ H]thymidine incorporation). ED50 ≤ 5 ng/ml.
Molecular Mass	Recombinant human Activin A is a 27.4 kDa disulphide-linked homodimers of two beta-A chains, each containing 116 amino residues.
AA Sequence	HHHHHHGLECDGKVNICKKQFFVSKDIGWNDWIIAPSGYHANYCEGECPSHIAG TSGSSLSFHSTVINHYRMR GHSPFANLKSCCVPTKLRPMSMLYYDDGQNIKKDIQN MIVEECGCS
Endotoxin	< 0.04 EU/ug protein (LAL)>
Purity	>97% by SDS-PAGE gel
Applications	Cell culture, Western blot
Storage	This lyophilized preparation is stable at 2-8o C for short term, long storage it should be kept at -20oC. Reconstituted protein should be stored in working aliquots at -20°C and it is recommended to add a carrier protein (0.1% HSA or BSA). Repeated freezing and thawing is not recommended.
Reconstitution	Lyophilized protein should be reconstituted in water to a concentration of 50 ng/ul.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Due to the protein nature, dimmers and multimers may be observed. Upon reconstitution, It can be stored in working aliquots at -20°C for future use. Optimal reconstitution please follow batch Quality Control sheet instructions.

GENE INFORMATION

Gene Name [INHBA inhibin, beta A \[Homo sapiens \]](#)

Official Symbol INHBA

Synonyms

INHBA; inhibin, beta A; inhibin, beta A (activin A, activin AB alpha polypeptide); inhibin beta A chain; Inhibin, beta-1; activin beta-A chain; FSH-releasing protein; erythroid differentiation factor; erythroid differentiation protein; follicle-stimulating hormone-releasing protein; EDF; FRP;

Gene ID [3624](#)

mRNA Refseq [NM_002192](#)

Protein Refseq [NP_002183](#)

MIM [147290](#)

UniProt ID [P08476](#)

Chromosome Location 7p15-p13

Pathway

ALK1 signaling events, organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Glycoprotein hormones, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Metabolism, organism-specific biosystem; Metabolism of amino acids and derivatives, organism-specific biosystem; Peptide hormone biosynthesis, organism-specific biosystem;

Function

cytokine activity; follistatin binding; growth factor activity; hormone activity; identical protein binding; protein binding; protein heterodimerization activity; type II activin receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA