

Recombinant Human INHBB, Animal Free

Cat. No. INHBB-138H Lot. No. (See product label)

SPECIFICATION

Product Overview	<p>Recombinant human Activin B is a 14 KDa betaB single chain, containing 123 amino residues. Human recombinant protein expressed in <i>Nicotiana benthamiana</i>. Recombinant human Activin B contains a 10-His-tag at the N-terminal end, is produced by transient expression in non-transgenic plants and is purified by sequential chromatography (FPLC). This product contains no animal-derived components or impurities. Animal free product.</p>
Species	<p>Human</p>
Source	<p><i>Nicotiana Benthamiana</i></p>
ProteinLength	<p>123 a.a.</p>
Description	<p>Activins are homodimers or heterodimers of the various beta subunit isoforms, belonging to the TGF-beta family. Mature Activin B has two chains of 123 amino acids residues (betaB-betaB). Activin exhibits a wide range of biological activities, including mesoderm induction, neural cell differentiation, bone remodelling, haematopoiesis, and reproductive physiology. Activins plays a key role in the production and regulation of hormones such as FSH, LH, GnRH and ACTH. Inhibins/Activins are protein that are formed by the dimerization of two subunits, i. e. an α (alpha) with either beta A (betaA) - Inhibin A or beta B (betaB) - Inhibin B. The subunits betaA and betaB can also form homodimers or heterodimers called Activin: Activin A (betaA -betaA), Activin B (betaB -betaB) and Activin AB (betaA -betaB). The Activin gene family comprises the additional, but poorly characterized members" Activin betaC (betaC), beta D</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

(betaD) and beta E (betaE). As with other members of the super-family, Activins interact with two types of cell surface trans-membrane receptors (Types I and II) which have intrinsic serine/threonine kinase activities in their cytoplasmic domains, Activin type 1 receptors, ACVR1, ACVR1B, ACVR1C and Activin type 2 receptors, ACVR2A, ACVR2B.

Form Lyophilized from Tris HCl 0.05M buffer at pH 7.4.

Molecular Mass Recombinant human Activin B is a 14 KDa betaB single chain, containing 123 amino residues.

AA Sequence HHHHHHHHHHGLECDGRTNLCCRQQFFIDFRLIGWNDWIIAPTGYGNYCEGSCPA
YLAGVPGSASSFHTAVVNQ YRMRGLNPGTVNSCCIPTKLSTMSMLYFDDEYNIVKR
DVPNMIVEECG

Endotoxin < 0.04 EU/ug protein (LAL)

Purity >97% by SDS-PAGE gel

Applications Western blot, Immunogen

Storage This lyophilized preparation is stable at 2-8o C for short term, long storage it should be kept at -20oC. Reconstituted protein should be stored in working aliquots at -20°C and it is recommended to add a carrier protein (0.1% HSA or BSA). Repeated freezing and thawing is not recommended.

Reconstitution Lyophilized protein should be reconstituted in water to a concentration of 50 ng/ul. Due to the protein nature, dimmers and multimers may be observed. Upon reconstitution, It can be stored in working aliquots at -20°C for future use. Optimal reconstitution please follow batch Quality Control sheet instructions.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GENE INFORMATION

Gene Name	INHBB inhibin, beta B [Homo sapiens]
Official Symbol	INHBB
Synonyms	INHBB; inhibin, beta B; inhibin, beta B (activin AB beta polypeptide); inhibin beta B chain; Inhibin, beta-2; activin beta-B chain; activin AB beta polypeptide; MGC157939;
Gene ID	3625
mRNA Refseq	NM_002193
Protein Refseq	NP_002184
MIM	147390
UniProt ID	P09529
Chromosome Location	2cen-q13
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Glycoprotein hormones, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of amino acids and derivatives, organism-specific biosystem; Peptide hormone biosynthesis, organism-specific biosystem; TGF-beta signaling pathway, organism-specific biosystem;
Function	cytokine activity; growth factor activity; hormone activity; host cell surface receptor binding; protein binding; protein homodimerization activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA