

Active Recombinant Human Insulin-Like 3 (Leydig cell)

Cat. No. INSL3-384H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human INSL3 protein was expressed in E. coli. Ala106-Tyr131(A chain) & Leu21-Ala55.
Species	Human
Source	E.coli
Protein Length	106-131;21-55 a.a.
Description	INSL3 is a member of the insulin/relaxin superfamily. INSL3 is a secreted protein produced in high amounts by testicular Leydig cells, and in lower amounts by ovarian follicular theca interna cells. As with other insulin/relaxin superfamily members, human INSL3 is synthesized as a prohormone. The Prepro INSL3 precursor is 131 amino acids (aa) in length. It is processed by a prohormone convertase to remove a 20 aa signal sequence and a 47 aa internal pro-, or C-peptide that separates a 35 aa N-terminal B chain from a 26 aa C-terminal A chain.
Form	Lyophilized from a 0.2 um filtered solution in PBS.
N-terminal Sequence	Ala106 (A chain) & Leu21 (B chain)
Molecular Weight	2.8 kDa (A chain), 3.3 kDa (B chain)
Activity	Measured by its ability to induce cAMP accumulation in human RXFP2 transfected HEK293 human embryonic kidney cells. The ED50 for this effect is typically 12.5-50

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	pg/mL.
Endotoxin Level	<1.0EU per 1µg of the protein by the LAL method.
Purity	>95%,by SDS-PAGE under reducing conditions and visualized by silver stain.
Reconstitution	Reconstituteat 100µg/mL in sterile PBS.
Storage	Use amannual defrost freezer and avoid repeated freezethaw cycles. 12 months from date of receipt, -20 to -70°C as supplied.1 month, 2 to 8°C under sterile conditions after reconstitution. 3 months,-20 to -70°C under sterile conditions after reconstitution.


GENE INFORMATION

Gene Name	INSL3 insulin-like 3 (Leydigcell) [Homo sapiens]
Official Symbol	INSL3
Synonyms	INSL3; insulin-like3 (Leydig cell); RLF; RLNL; insulin-like 3; ley-I-L; relaxin-likefactor b; leydig insulin-like peptide; leydig insulin -like hormone
Gene ID	3640
mRNA Refseq	NM_005543
Protein Refseq	NP_005534
MIM	146738
UniProt ID	P51460

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	19p13.2-p12
Pathway	Class A/1 (Rhodopsin-like receptors); Cytokine Signaling in Immune system; G alpha (s) signalling events; GPCR downstream signaling; GPCR ligand binding; Immune System; Interleukin receptor SHC signaling; Interleukin-2 signaling; Interleukin-3, 5 and GM-CSF signaling; Interleukin-7 signaling; Peptide ligand-binding receptors; Relaxin receptors; Signal Transduction; Signaling by GPCR; Signaling by Interleukins
Function	G-protein coupled receptor binding; hormone activity; insulin receptor binding; receptor binding
PDB rendering based on 2h8b.	

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA