

Recombinant Human INSR cell lysate

Cat. No. INSR-474HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human Insulin Receptor / INSR / CD220 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[INSR insulin receptor \[Homo sapiens \]](#)

Official Symbol

INSR

Synonyms

INSR; insulin receptor; CD220; IR; HHF5;

Gene ID

[3643](#)

mRNA Refseq

[NM_000208](#)

Protein Refseq

[NP_000199](#)

MIM

[147670](#)

UniProt ID

[P06213](#)

Chromosome Location

19p13.3-p13.2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway


Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Aldosterone-regulated sodium reabsorption, organism-specific biosystem; Aldosterone-regulated sodium reabsorption, conserved biosystem; IRS activation, organism-specific biosystem; IRS-mediated signalling, organism-specific biosystem; IRS-related events, organism-specific biosystem;

Function

3-phosphoinositide-dependent protein kinase binding; ATP binding; GTP binding; PTB domain binding; SH2 domain binding; insulin binding; insulin binding; insulin binding; insulin receptor substrate binding; insulin-activated receptor activity; insulin-like growth factor I binding; insulin-like growth factor II binding; insulin-like growth factor receptor binding; lipoic acid binding; nucleotide binding; phosphatidylinositol 3-kinase binding; protein binding; protein complex binding; protein domain specific binding; protein phosphatase binding; protein tyrosine kinase activity; protein tyrosine kinase activity; protein tyrosine kinase activity; receptor activity; receptor signaling protein tyrosine kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA