

# Active Recombinant Human Integrin Alpha 5 Beta 3 Heterodimer Protein

**Cat. No.** ITGAVB3-101H    **Lot. No.** (See product label)

## SPECIFICATION

<b>Product Overview</b>	Recombinant human Integrin alpha 5 beta 3 heterodimer protein (Human ITGA5 & ITGB3) Phe 42 - Tyr 995 (Integrin alpha 5, NP_002196) and Gly 27 - Asp 718 (Integrin beta 3, AAI27667) was co-expressed in human 293 cells (HEK293).
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	42-995;27-718 a.a.
<b>Description</b>	The product of this gene belongs to the integrin alpha chain family. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This gene encodes the integrin alpha 5 chain. Alpha chain 5 undergoes post-translational cleavage in the extracellular domain to yield disulfide-linked light and heavy chains that join with beta 1 to form a fibronectin receptor. In addition to adhesion, integrins are known to participate in cell-surface mediated signalling.
<b>Molecular Mass</b>	Subunit Integrin alpha 5 (ITGA5) is fused with Strep II-tag at the C-terminus and subunit Integrin beta 3 (ITGB3) is fused with his-tag at the C-terminus. The heterodimer Human ITGA5 & ITGB3, co-expressed by ITGA5 plasmid and ITGB3 plasmid, has a calculated MW of 105.4 kDa ( alpha 5 subunit) and 77.3 kDa (beta 3 subunit). The predicted N-terminus is Phe42 (Integrin alpha 5) and Gly 27 (Integrin beta 3) . DTT-reduced Protein migrates as 105-150 kDa due to glycosylation.

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Formulation</b>	Lyophilized from 0.22 µm filtered solution in 20 mM Tris, pH 7.5 with 150 mM NaCl and 0.5 mM CaCl <sub>2</sub> . Normally Mannitol or Trehalose are added as protectants before lyophilization.
<b>Purity</b>	>90% as determined by SDS-PAGE.
<b>Bioactivity</b>	Measured by its ability in functional ELISA. Immobilized recombinant human Vitronectin protein at 1 µg/ml can bind Recombinant human Integrin alpha 5 beta 3 heterodimer protein with apparent KD <8 nM.
<b>Endotoxin</b>	Less than 1.0 EU per µg of the Human ITGA5 & ITGB3 by the LAL method.
<b>Storage</b>	-20°C to -70°C for 3 months under sterile conditions after reconstitution. Avoid repeated freezethaw cycles.
<b>OfficialSymbol</b>	ITGA5
<b>GENE INFORMATION</b>	
<b>Gene Name</b>	<a href="#">ITGA5 integrin, alpha 5 (fibronectin receptor, alpha polypeptide) [ Homo sapiens ]</a>
<b>Synonyms</b>	ITGA5; integrin, alpha 5 (fibronectin receptor, alpha polypeptide); FNRA; CD49e; VLA5A; integrin alpha-5; VLA-5; integrin alpha-F; CD49 antigen-like family member E; fibronectin receptor subunit alpha; fibronectin receptor, alpha subunit; very late activation protein 5, alpha subunit; Integrin alpha-5; Fibronectin receptor subunit alpha; Integrin alpha-F
<b>Gene ID</b>	<a href="#">3678</a>
<b>mRNA Refseq</b>	<a href="#">NM_002205</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Protein Refseq</b>	NP_002196
<b>MIM</b>	135620
<b>UniProt ID</b>	P08648
<b>Chromosome Location</b>	12q11-q13
<b>Pathway</b>	Angiopoietin receptor Tie2-mediated signaling; Arrhythmogenic right ventricular cardiomyopathy (ARVC); Axon guidance; Bacterial invasion of epithelial cells; Cell surface interactions at the vascular wall; Dilated cardiomyopathy; ECM-receptor interaction; Focal Adhesion; Focal adhesion; Hematopoietic cell lineage; Hemostasis; Integrin cell surface interactions; Integrin-mediated cell adhesion; L1CAM interactions; Pertussis
<b>Function</b>	protein binding; metal ion binding; platelet-derived growth factor receptor binding; vascular endothelial growth factor receptor 2 binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA