

## Native Goat IgG Fab fragment

Cat. No. IgG-150G Lot. No. (See product label)


### SPECIFICATION

<b>Species</b>	Goat
<b>Source</b>	Goat Serum
<b>ProteinLength</b>	Full length native Goat IgG Fab fragment
<b>Description</b>	The fragment antigen-binding (Fab fragment) is a region on an antibody that binds to antigens. It is composed of one constant and one variable domain of each of the heavy and the light chain. These domains shape the paratope — the antigen-binding site — at the amino terminal end of the monomer. The two variable domains bind the epitope on their specific antigens.
<b>Form</b>	Liquid
<b>Purity</b>	>95% (SDS-PAGE)
<b>Applications</b>	Western blotting, ELISA, etc
<b>Storage</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
<b>Storage Buffer</b>	PBS pH 7.4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA