

## Recombinant Mouse Interleukin 13

Cat. No. II13-68M Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Mouse Interleukin-13 produced in <i>E. Coli</i> is a single, non-glycosylated polypeptide chain containing 110 amino acids and having a molecular mass of 12127 Dalton. The IL-13 is purified by proprietary chromatographic techniques.
<b>Species</b>	Mouse
<b>Source</b>	E.coli
<b>Description</b>	IL13 is an immunoregulatory cytokine produced primarily by activated Th2 cells. IL-13 is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen-induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4.
<b>Amino Acid Sequence</b>	The sequence of the first five N-terminal amino acids was determined and found to be Met-Pro-Val-Pro-Arg.
<b>Physical Appearance</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Purity</b>	Greater than 98% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Formulation</b>	The protein (1mg/ml) was lyophilized with no additives.
<b>Solubility</b>	It is recommended to reconstitute the lyophilized Interleukin 13 in sterile 18MΩ-cm H <sub>2</sub> O not less than 100g/ml, which can then be further diluted to other aqueous solutions.
<b>Biological Activity</b>	The ED <sub>50</sub> was determined by the dose dependent proliferation of TF-1 cells and was found to be < 4ng/ml, corresponding to a specific activity of >2.5 x 10 <sup>5</sup> .
<b>Protein Content</b>	Protein quantitation was carried out by two independent methods: 1. UV spectroscopy at 280 nm using the absorbency value of 0.69 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics). 2. Analysis by RP-HPLC, using a calibrated solution of IL-13 as a Reference Standard.
<b>Storage</b>	Lyophilized Interleukin-13 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL13 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

## GENE INFORMATION

<b>Gene Name</b>	Il13 interleukin 13 [ <i>Mus musculus</i> ]
<b>Synonyms</b>	Il13; interleukin 13; Il-13; OTTMUSP00000005816
<b>Gene ID</b>	16163
<b>mRNA Refseq</b>	NM_008355


 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Protein Refseq</b>	<a href="#">NP_032381</a>
<b>UniProt ID</b>	<a href="#">P20109</a>
<b>Chromosome Location</b>	11 B1.3; 11 29.0 cM
<b>Pathway</b>	Asthma; Cytokine-cytokine receptor interaction; Fc epsilon RI signaling pathway; Jak-STAT signaling pathway
<b>Function</b>	cytokine activity;cytokine receptor binding; interleukin-13 receptor binding

**PDB rendering based on 1ga3.**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA