

## Recombinant Rat Interleukin 17F

Cat. No. II17f-204R Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Rat Interleukin-17F produced in <i>E. Coli</i> is a homodimeric, non-glycosylated polypeptide chain containing 145 amino acids and having a molecular mass of 35.5 kDa. The IL-25 Rat is purified by proprietary chromatographic techniques.
<b>Species</b>	Rat
<b>Source</b>	E.coli
<b>Description</b>	<p>IL-17F having an accession number of Q96PD4 is a cytokine that shares sequence similarity with IL17. IL-17F is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM-CSF. IL-17F inhibits the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1. IL-17F induces stromal cells to produce proinflammatory and hematopoietic cytokines. Intestinal IL17F gene expression is increased in active CD. IL-17A and IL-17F alleles influence the susceptibility to and pathophysiological features of ulcerative colitis independently. IL-17F and MIF gene polymorphisms are significantly associated with the development of functional dyspepsia. The initiation of IL-17F/IL-17R signaling pathway requires the receptor ubiquitination by TRAF6. IL-17F induces expression of IFN-gamma-inducible protein 10 (IP-10) by activating Raf1-mitogen-activated protein kinase 1/2-extracellular-regulated kinase 1/2-p90 ribosomal S6 kinase-cyclic AMP response element-binding protein signaling pathway.</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Amino Acid Sequence</b>	MARRNPKVGL SALQKAGNCP PLEDNSVRVD IRIFNQNGGI SVPRDFQNRS SSPWDYNITR DPDRFPSEIA EAQCRHSGCI NAQQQEDGSM NSVPIQQEIL VLRREPQGCS NSFRLKMLI KVGCTCVTPI VHHA.
<b>Physical Appearance</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Purity</b>	Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
<b>Formulation</b>	IL17F Rat was lyophilized from a concentrated (1mg/ml) solution containing no additives.
<b>Solubility</b>	It is recommended to reconstitute the lyophilized Interleukin Rat IL17F in sterile 18MΩ-cm H <sub>2</sub> O not less than 100g/ml, which can then be further diluted to other aqueous solutions.
<b>Stability</b>	Lyophilized Interleukin Rat 17F although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL17F should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">Il17f interleukin 17F [ Rattus norvegicus ]</a>
<b>Synonyms</b>	Il17f; interleukin 17F; Il17f_predicted; LOC301291; MGC114402; interleukin 17F (predicted)
<b>Gene ID</b>	<a href="#">301291</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



<b>mRNA Refseq</b>	NM_001015011
<b>Protein Refseq</b>	NP_001015011
<b>UniProt ID</b>	Q5BJ95
<b>Chromosome Location</b>	9q13
<b>Function</b>	cytokine activity;cytokine binding; protein homodimerization activity

Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA