

## Active Recombinant Human IL23A, HIgG1 Fc-tagged, mutant

Cat. No. IL23A-001H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	The human IL-23 complex composed of the IL23a subunit(NP_057668.1) (Met1-Pro189) and the IL12b subunit(NP_002178.2)(Ile23-Ser328) is fused through a polypeptide linker to the Fc region of mutant human IgG1was expressed in CHO cell.
<b>Species</b>	Human
<b>Source</b>	CHO
<b>ProteinLength</b>	1-189;23-328 a.a.
<b>Description</b>	This gene encodes a subunit of the heterodimeric cytokine interleukin 23 (IL23). IL23 is composed of this proteinand the p40 subunit of interleukin 12 (IL12B). The receptor of IL23 is formed by the beta 1 subunit of IL12(IL12RB1) and an IL23 specific subunit, IL23R. Both IL23 and IL12 can activate the transcription activator STAT4,and stimulate the production of interferon-gamma (IFNG). In contrast to IL12, which acts mainly on naive CD4(+) Tcells, IL23 preferentially acts on memory CD4(+) T cells
<b>Form</b>	Lyophilized from 0.2µm-filtered solution in PBS.
<b>Bio-activity</b>	Measured by its binding ability in a functional ELISA.
<b>Molecular Mass</b>	81KDa (monomer)
<b>AA Sequence</b>	Met1-Pro189 .

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Endotoxin</b>	<0.06 eu/μg="" as="" determined="" by="" la="">
<b>Purity</b>	>98%, by SDS-PAGE under reducing conditions
<b>Stability</b>	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
<b>Reconstitution</b>	Reconstitute at 100μg/ml in sterile PBS.
<b>Warning</b>	Avoid freeze/thaw cycles.

## GENE INFORMATION

<b>Gene Name</b>	IL23A interleukin 23, alpha subunit p19 [ Homo sapiens ]
<b>Official Symbol</b>	IL23A
<b>Synonyms</b>	IL23A; interleukin 23, alpha subunit p19; interleukin-23 subunit alpha; IL 23; IL 23A; IL23P19; interleukin six; G CSF related factor; P19; SGRF; IL-23-A; IL-23p19; IL-23 subunit alpha; interleukin 23 p19 subunit; interleukin-23 subunit p19; JKA3 induced upon T-cell activation; interleukin-six, G-CSF related factor; IL-23; IL-23A; MGC79388;
<b>Gene ID</b>	51561
<b>mRNA Refseq</b>	NM_016584
<b>Protein Refseq</b>	NP_057668
<b>MIM</b>	605580

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>UniProt ID</b>	Q9NPF7
<b>Chromosome Location</b>	12q13.13
<b>Pathway</b>	ATF-2 transcription factor network, organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; IL23-mediated signaling events, organism-specific biosystem; Jak-STAT signaling pathway, organism-specific biosystem; Jak-STAT signaling pathway, conserved biosystem; Pertussis, organism-specific biosystem;
<b>Function</b>	cytokine activity; contributes_to interleukin-23 receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA