

## Recombinant Mouse Insig2 Protein, Myc/DDK-tagged

**Cat. No.** Insig2-3552M    **Lot. No.** (See product label)

### SPECIFICATION

**Product Overview**      Purified recombinant protein of mouse full-length insulin induced gene 2 (Insig2), with C-terminal MYC/DDK tag, expressed in HEK293T cells.

**Species**                      Mouse

**Source**                        HEK293

**Description**

Oxysterol-binding protein that mediates feedback control of cholesterol synthesis by controlling both endoplasmic reticulum to Golgi transport of SCAP and degradation of HMGCR. Acts as a negative regulator of cholesterol biosynthesis by mediating the retention of the SCAP-SREBP complex in the endoplasmic reticulum, thereby blocking the processing of sterol regulatory element-binding proteins (SREBPs) SREBF1/SREBP1 and SREBF2/SREBP2. Binds oxysterol, including 22-hydroxycholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol and 27-hydroxycholesterol, regulating interaction with SCAP and retention of the SCAP-SREBP complex in the endoplasmic reticulum. In presence of oxysterol, interacts with SCAP, retaining the SCAP-SREBP complex in the endoplasmic reticulum, thereby preventing SCAP from escorting SREBF1/SREBP1 and SREBF2/SREBP2 to the Golgi. Sterol deprivation or phosphorylation by PCK1 reduce oxysterol-binding, disrupting the interaction between INSIG2 and SCAP, thereby promoting Golgi transport of the SCAP-SREBP complex, followed by processing and nuclear translocation of SREBF1/SREBP1 and SREBF2/SREBP2. Also regulates cholesterol synthesis by regulating degradation of HMGCR: initiates the sterol-mediated ubiquitin-mediated endoplasmic reticulum-associated degradation (ERAD) of HMGCR

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

via recruitment of the reductase to the ubiquitin ligase RNF139.

<b>Molecular Mass</b>	24.9 kDa
<b>Purity</b>	> 80% as determined by SDS-PAGE and Coomassie blue staining
<b>Stability</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>Storage</b>	Store at -80 centigrade after receiving vials.
<b>Concentration</b>	>50 µg/mL as determined by microplate BCA method
<b>Storage Buffer</b>	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.

## GENE INFORMATION

<b>Gene Name</b>	Insig2 insulin induced gene 2 [ <i>Mus musculus</i> (house mouse) ]
<b>Official Symbol</b>	<a href="#">Insig2</a>
<b>Synonyms</b>	Insig2; insulin induced gene 2; Ins; Insig-2; 2900053I11Rik; C730043J18Rik; insulin-induced gene 2 protein; INSIG-2 membrane protein
<b>Gene ID</b>	<a href="#">72999</a>
<b>mRNA Refseq</b>	<a href="#">NM_178082</a>
<b>Protein Refseq</b>	<a href="#">NP_835183</a>
<b>UniProt ID</b>	<a href="#">Q91WG1</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA