

Recombinant Human JAK2 Protein, His-tagged

Cat. No. JAK2-01H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human Jak2 protein (1014-1132aa), fused to His-tag at N-terminus, was expressed in E. coli
Species	Human
Source	E.coli
ProteinLength	1014-1132 a.a.
Description	<p>This gene encodes a non-receptor tyrosine kinase that plays a central role in cytokine and growth factor signalling. The primary isoform of this protein has an N-terminal FERM domain that is required for erythropoietin receptor association, an SH2 domain that binds STAT transcription factors, a pseudokinase domain and a C-terminal tyrosine kinase domain. Cytokine binding induces autophosphorylation and activation of this kinase. This kinase then recruits and phosphorylates signal transducer and activator of transcription (STAT) proteins. Growth factors like TGF-beta 1 also induce phosphorylation and activation of this kinase and translocation of downstream STAT proteins to the nucleus where they influence gene transcription. Mutations in this gene are associated with numerous inflammatory diseases and malignancies. This gene is a downstream target of the pleiotropic cytokine IL6 that is produced by B cells, T cells, dendritic cells and macrophages to produce an immune response or inflammation. Disregulation of the IL6/JAK2/STAT3 signalling pathways produces increased cellular proliferation and myeloproliferative neoplasms of hematopoietic stem cells. A nonsynonymous mutation in the pseudokinase domain of this gene</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

disrupts the domains inhibitory effect and results in constitutive tyrosine phosphorylation activity and hypersensitivity to cytokine signalling. This gene and the IL6/JAK2/STAT3 signalling pathway is a therapeutic target for the treatment of excessive inflammatory responses to viral infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Form	Liquid
Molecular Mass	18.1 kDa (157aa)
AA Sequence	MRGSHHHHHHGMASMTGGQQMGRDLYDDDDKDRWGSHEMGEPIFWYAPESLTSKFSVASDVWSFGVVLVELFTYIEKSKSPPAEFMRMIGNDKQGQMIVFHLLIELLKNNGRLRPRDGCPCDEIYMIMTECWNNNVNQRPSFRDLALRVDQIRDNMAG
Purity	> 95% as analyzed by SDS-PAGE
Applications	SDS-PAGE, Denatured
Storage	Can be stored at 2 to 8 centigrade for 1 week. For long term storage, aliquot and store at -20 to -80 centigrade. Avoid repeated freezing and thawing cycles.
Concentration	1 mg/mL (determined by Bradford assay)
Storage Buffer	20mM Tris-HCl buffer (pH 8.0) containing, 10% glycerol, 0.4M urea

GENE INFORMATION

Gene Name	JAK2 Janus kinase 2 [Homo sapiens (human)]
Official Symbol	JAK2


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	JAK2; Janus kinase 2; JTK10; tyrosine-protein kinase JAK2; JAK-2; Janus kinase 2 (a protein tyrosine kinase); EC 2.7.10.2
Gene ID	3717
mRNA Refseq	NM_004972
Protein Refseq	NP_004963
MIM	147796
UniProt ID	O60674

SDS-PAGE



3 µg by SDS-PAGE under reducing condition and visualized by coomassie blue stain.