

Active Recombinant Human JAK2 (JH1 domain) Protein, His-tagged

Cat. No. JAK2-23H Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant human JAK2 (Janus Kinase 2) encompassing amino acids 808-1132 (end), containing the tyrosine kinase domain JH1. This construct contains an N-terminal His-tag (6xHis). The protein was affinity purified and is kinase active.

Species Human

Source Insect Cells

ProteinLength 808-1132

Description This gene encodes a non-receptor tyrosine kinase that plays a central role in cytokine and growth factor signalling. The primary isoform of this protein has an N-terminal FERM domain that is required for erythropoietin receptor association, an SH2 domain that binds STAT transcription factors, a pseudokinase domain and a C-terminal tyrosine kinase domain. Cytokine binding induces autophosphorylation and activation of this kinase. This kinase then recruits and phosphorylates signal transducer and activator of transcription (STAT) proteins. Growth factors like TGF-beta 1 also induce phosphorylation and activation of this kinase and translocation of downstream STAT proteins to the nucleus where they influence gene transcription. Mutations in this gene are associated with numerous inflammatory diseases and malignancies. This gene is a downstream target of the pleiotropic cytokine IL6 that is produced by B cells, T cells, dendritic cells and macrophages to produce an immune response or inflammation. Disregulation of the IL6/JAK2/STAT3 signalling pathways produces

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

increased cellular proliferation and myeloproliferative neoplasms of hematopoietic stem cells. A nonsynonymous mutation in the pseudokinase domain of this gene disrupts the domains inhibitory effect and results in constitutive tyrosine phosphorylation activity and hypersensitivity to cytokine signalling. This gene and the IL6/JAK2/STAT3 signalling pathway is a therapeutic target for the treatment of excessive inflammatory responses to viral infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms.


Form	Aqueous buffer solution
Bio-activity	124 pmol/min/μg.
Molecular Mass	43 kDa
Purity	≥ 70%
Applications	Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.
Usage	Assay was done according to JAK2 (Janus Kinase 2) Assay Kit with 20 μM ATP instead of 10 μM ATP and various amounts of JAK2 enzyme.
Storage	At least 6 months at –80 centigrade.
Storage Buffer	40 mM Tris-HCl, pH 8.0, 110 mM NaCl, 2.2 mM KCl, 3 mM DTT, 0.04% Tween-20 and 20% Glycerol

GENE INFORMATION


Gene Name	JAK2 Janus kinase 2 [Homo sapiens (human)]
Official Symbol	JAK2

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	JAK2; Janus kinase 2; JTK10; tyrosine-protein kinase JAK2; JAK-2; Janus kinase 2 (a protein tyrosine kinase); EC 2.7.10.2; jak; jh1
Gene ID	3717
mRNA Refseq	NM_004972
Protein Refseq	NP_004963
MIM	147796
UniProt ID	O60674




Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA