

Recombinant Human JAK2 (V617F) Protein, FLAG/Avi-tagged, Biotin-labeled

Cat. No. JAK2-27H Lot. No. (See product label)

SPECIFICATION

Product Overview	Full length human JAK2, also known as Janus kinase-2, Just Another Kinase 2, with V617F mutation, GenBank Accession No. NM_004972, a.a. 1-1132 (end) (full length), with C-terminal FLAGAvi-Tag, co-expressed with PTP1B in HEK293 expression system. This protein is less than 10% phosphorylated. This protein is enzymatically Biotin-labeled using AviTag technology.
Species	Human
Source	HEK293
ProteinLength	1-1132
Description	This gene encodes a non-receptor tyrosine kinase that plays a central role in cytokine and growth factor signalling. The primary isoform of this protein has an N-terminal FERM domain that is required for erythropoietin receptor association, an SH2 domain that binds STAT transcription factors, a pseudokinase domain and a C-terminal tyrosine kinase domain. Cytokine binding induces autophosphorylation and activation of this kinase. This kinase then recruits and phosphorylates signal transducer and activator of transcription (STAT) proteins. Growth factors like TGF-beta 1 also induce phosphorylation and activation of this kinase and translocation of downstream STAT proteins to the nucleus where they influence gene transcription. Mutations in this gene are associated with numerous inflammatory diseases and malignancies. This gene is a downstream target of the pleiotropic cytokine IL6 that is produced by B

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cells, T cells, dendritic cells and macrophages to produce an immune response or inflammation. Disregulation of the IL6/JAK2/STAT3 signalling pathways produces increased cellular proliferation and myeloproliferative neoplasms of hematopoietic stem cells. A nonsynonymous mutation in the pseudokinase domain of this gene disrupts the domains inhibitory effect and results in constitutive tyrosine phosphorylation activity and hypersensitivity to cytokine signalling. This gene and the IL6/JAK2/STAT3 signalling pathway is a therapeutic target for the treatment of excessive inflammatory responses to viral infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Form Aqueous buffer solution

Molecular Mass 134 kDa

Storage At least 6 months at -80 centigrade.

Storage Buffer 40 mM Tris-HCl, pH 8.0, 110 mM NaCl, 2.2 mM KCl, 0.04% Tween-20, 100µg/mL FLAG peptide and 20% glycerol.

Shipping At -80 centigrade

Conjugation Biotin

GENE INFORMATION

Gene Name JAK2 Janus kinase 2 [Homo sapiens (human)]

Official Symbol JAK2

Synonyms JAK2; Janus kinase 2; JTK10; tyrosine-protein kinase JAK2; JAK-2; Janus kinase 2 (a protein tyrosine kinase); EC 2.7.10.2; jak; jh1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 3717


mRNA Refseq NM_004972

Protein Refseq NP_004963

MIM 147796

UniProt ID O60674

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA