

## Recombinant Human JAK2 (V617F) (JH2 Domain) Protein, His/Avi-tagged, Biotin-labeled

Cat. No. JAK2-30H Lot. No. (See product label)

### SPECIFICATION

**Product Overview** Human JAK2, also known as Janus kinase-2, Just Another Kinase 2, with V617F mutation, GenBank Accession No. NM\_004972, a.a. 513-827, with C-terminal His-Avi-Tag, and expressed in an Sf9 expression system. This protein is enzymatically Biotin-labeled using Avi-Tag technology.

**Species** Human

**Source** Insect Cells

**ProteinLength** 513-827

**Description** This gene encodes a non-receptor tyrosine kinase that plays a central role in cytokine and growth factor signalling. The primary isoform of this protein has an N-terminal FERM domain that is required for erythropoietin receptor association, an SH2 domain that binds STAT transcription factors, a pseudokinase domain and a C-terminal tyrosine kinase domain. Cytokine binding induces autophosphorylation and activation of this kinase. This kinase then recruits and phosphorylates signal transducer and activator of transcription (STAT) proteins. Growth factors like TGF-beta 1 also induce phosphorylation and activation of this kinase and translocation of downstream STAT proteins to the nucleus where they influence gene transcription. Mutations in this gene are associated with numerous inflammatory diseases and malignancies. This gene is a downstream target of the pleiotropic cytokine IL6 that is produced by B cells, T cells, dendritic cells and macrophages to produce an immune response or

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

inflammation. Disregulation of the IL6/JAK2/STAT3 signalling pathways produces increased cellular proliferation and myeloproliferative neoplasms of hematopoietic stem cells. A nonsynonymous mutation in the pseudokinase domain of this gene disrupts the domains inhibitory effect and results in constitutive tyrosine phosphorylation activity and hypersensitivity to cytokine signalling. This gene and the IL6/JAK2/STAT3 signalling pathway is a therapeutic target for the treatment of excessive inflammatory responses to viral infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

**Form** Aqueous buffer solution

**Molecular Mass** 39 kDa

**Storage** At least 6 months at -80 centigrade.

**Storage Buffer** 40 mM Tris-HCl, pH 8.0, 110 mM NaCl, 2.2 mM KCl, 0.04% Tween-20, 500 mM Imidazole and 20% glycerol.

**Shipping** At -80 centigrade

**Conjugation** Biotin

## GENE INFORMATION

**Gene Name** JAK2 Janus kinase 2 [ Homo sapiens (human) ]

**Official Symbol** JAK2

**Synonyms** JAK2; Janus kinase 2; JTK10; tyrosine-protein kinase JAK2; JAK-2; Janus kinase 2 (a protein tyrosine kinase); EC 2.7.10.2; jak; jh1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Gene ID	3717
mRNA Refseq	NM_004972
Protein Refseq	NP_004963
MIM	147796
UniProt ID	O60674

**SDS-PAGE**

**4-20% SDS-PAGE**  
**Coomassie staining**

**Lane 1:**  
2µg JAK2(V617F)  
(513-827-  
His-Avi-Biotin)

**Lane 2:**  
Protein Marker  
**MW:** 39 kDa  
**Purity:** ≥58%



1 2

— 250 kDa  
— 150 kDa  
— 100 kDa  
— 75 kDa  
— 50 kDa  
— 37 kDa  
— 25 kDa  
— 20 kDa  
— 15 kDa  
— 10 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA