

## Recombinant Human JAK2 (V617F, W659A, W777A, F794H) Mutation Protein, His/Avi-Tag, Biotin-Labeled

Cat. No. JAK2-37H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant human JAK2 (Janus Kinase 2) encompassing amino acids 536-812, containing the pseudo-kinase domain JH2 with mutations V617F, W659A, W777A, and F794H. This construct contains a C-terminal His-tag (6xHis) followed by an Avi-Tag™. The protein was enzymatically biotinylated using the Avi-Tag™ and affinity purified. This protein is enzymatically biotinylated using Avi-Tag™ technology. Biotinylation confirmed to be > 90%.
<b>Species</b>	Human
<b>Source</b>	Insect Cells
<b>ProteinLength</b>	536-812
<b>Description</b>	This gene encodes a non-receptor tyrosine kinase that plays a central role in cytokine and growth factor signalling. The primary isoform of this protein has an N-terminal FERM domain that is required for erythropoietin receptor association, an SH2 domain that binds STAT transcription factors, a pseudokinase domain and a C-terminal tyrosine kinase domain. Cytokine binding induces autophosphorylation and activation of this kinase. This kinase then recruits and phosphorylates signal transducer and activator of transcription (STAT) proteins. Growth factors like TGF-beta 1 also induce phosphorylation and activation of this kinase and translocation of downstream STAT proteins to the nucleus where they influence gene transcription. Mutations in this gene are associated with numerous inflammatory diseases and malignancies. This

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

gene is a downstream target of the pleiotropic cytokine IL6 that is produced by B cells, T cells, dendritic cells and macrophages to produce an immune response or inflammation. Disregulation of the IL6/JAK2/STAT3 signalling pathways produces increased cellular proliferation and myeloproliferative neoplasms of hematopoietic stem cells. A nonsynonymous mutation in the pseudokinase domain of this gene disrupts the domains inhibitory effect and results in constitutive tyrosine phosphorylation activity and hypersensitivity to cytokine signalling. This gene and the IL6/JAK2/STAT3 signalling pathway is a therapeutic target for the treatment of excessive inflammatory responses to viral infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

<b>Form</b>	Aqueous buffer solution
<b>Molecular Mass</b>	35 kDa
<b>Purity</b>	≥ 90%
<b>Storage</b>	At least 6 months at –80 centigrade.
<b>Storage Buffer</b>	45 mM Tris-HCl, pH 8.0, 270 mM NaCl, 2 mM KCl, 20% glycerol, 0.5 mM TCEP
<b>Conjugation</b>	Biotin

## GENE INFORMATION



<b>Gene Name</b>	JAK2 Janus kinase 2 [ Homo sapiens (human) ]
<b>Official Symbol</b>	JAK2
<b>Synonyms</b>	JAK2; Janus kinase 2; JTK10; tyrosine-protein kinase JAK2; JAK-2; Janus kinase 2 (a protein tyrosine kinase); EC 2.7.10.2; jak; jh1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Gene ID</b>	3717
<b>mRNA Refseq</b>	NM_004972
<b>Protein Refseq</b>	NP_004963
<b>MIM</b>	147796
<b>UniProt ID</b>	O60674



Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA