

Recombinant Human KCTD9 293 Cell Lysate

Cat. No. KCTD9-5004HCL **Lot. No.** (See product label)

SPECIFICATION

| | |
|----------------------------|---|
| Species | Human |
| Source | HEK293 |
| Description | Antigen standard for potassium channel tetramerisation domain containing 9 (KCTD9) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection. |
| Components | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| Size | 0.1 mg |
| Storage Instruction | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment. |
| Applications | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

| | |
|----------------------------|--|
| Gene Name | KCTD9 potassium channel tetramerisation domain containing 9 [Homo sapiens] |
| Official Symbol | KCTD9 |
| Synonyms | KCTD9; potassium channel tetramerisation domain containing 9; BTB/POZ domain-containing protein KCTD9; FLJ20038; |
| Gene ID | 54793 |
| mRNA Refseq | NM_017634 |
| Protein Refseq | NP_060104 |
| UniProt ID | Q7L273 |
| Chromosome Location | 8p21.1 |
| Function | voltage-gated potassium channel activity; |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA