

Recombinant Human KDM3A 293 Cell Lysate

Cat. No. KDM3A-4997HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for lysine (K)-specific demethylase 3A (KDM3A), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	KDM3A lysine (K)-specific demethylase 3A [Homo sapiens]
Official Symbol	KDM3A
Synonyms	KDM3A; lysine (K)-specific demethylase 3A; JMJD1, JMJD1A, jumonji domain containing 1 , jumonji domain containing 1A; lysine-specific demethylase 3A; JHMD2A; jumonji C domain containing histone demethylase 2A; KIAA0742; TSGA; testis-specific protein A; jumonji domain containing 1; jumonji domain containing 1A; jumonji domain-containing protein 1A; jumonji C domain-containing histone demethylase 2A; jmjC domain-containing histone demethylation protein 2A; JMJD1; JHDM2A; JMJD1A; DKFZp686A24246; DKFZp686P07111;
Gene ID	55818
mRNA Refseq	NM_001146688
Protein Refseq	NP_001140160
MIM	611512
UniProt ID	Q9Y4C1
Chromosome Location	2p11.2
Pathway	Coregulation of Androgen receptor activity, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

androgen receptor binding; iron ion binding; metal ion binding; oxidoreductase activity; oxidoreductase activity, acting on single donors with incorporation of molecular oxygen, incorporation of two atoms of oxygen;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA