

Recombinant Mouse KDR cell lysate

Cat. No. KDR-1520MCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Mouse KDR / VEGFR2 / CD309 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	Kdr kinase insert domain protein receptor [Mus musculus]
Official Symbol	KDR
Synonyms	KDR; kinase insert domain protein receptor; vascular endothelial growth factor receptor 2; kinase NYK; VEGF receptor-2; fetal liver kinase 1; protein-tyrosine kinase receptor flk-1; vascular endothelial growth factor receptor-2; vascular endothelial growth factor receptor-3; vascular endothelial growth factor receptor- 2; soluble vascular endothelial growth factor receptor 2; Flk1; Ly73; Flk-1; Krd-1; VEGFR2; VEGFR-2; sVEGFR-2; 6130401C07;
Gene ID	16542
mRNA Refseq	NM_010612
Protein Refseq	NP_034742
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cytokine receptor interaction, conserved biosystem; Endocytosis, organism-specific biosystem; Endocytosis, conserved biosystem; Focal adhesion, organism-specific biosystem; Focal adhesion, conserved biosystem; Id Signaling Pathway, organism-specific biosystem;

Function

ATP binding; growth factor binding; growth factor binding; integrin binding; kinase activity; nucleotide binding; protein tyrosine kinase activity; receptor activity; transferase activity; vascular endothelial growth factor-activated receptor activity; vascular endothelial growth factor-activated receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA