

Active Recombinant Human Killer Cell Immunoglobulin-Like Receptor, Three Domains, Long Cytoplasmic Tail, 2, Fc-tagged

Cat. No. KIR3DL2-5669H **Lot. No.** (See product label)

SPECIFICATION

Product Overview Recombinant Human KIR3DL2 protein was expressed in Murine myeloma cell line with the Fc part of Human IgG1 Pro100-Lys330 (C-terminus) and Human KIR3DL2 Leu22-Leu339(N-terminus).

Species Human

Source Mammalian Cells

Protein Length 22-339 a.a.

Description KIR3DL2 is a type I transmembrane protein of the p70 family of killer cell Ig-like receptors (KIR). KIR are expressed on CD56 dim NK cells and T cell subsets where they participate in identifying normal and abnormal cells and regulating effector functions of the innate immune system. KIR are named for the number of Ig-like domains (2D or 3D) in the extracellular domain (ECD) and whether they have long or short (L, S) cytoplasmic tail. As with other inhibitory KIR, KIR3DL2 has two ITIM domains within its long tail (3). KIR3DL2 is diverse, with twelve alleles identified and as many as five single amino acid (aa) polymorphisms found in a single individual.

Form Lyophilized from a 0.2 µm filtered solution in PBS.

N-terminal Sequence Leu22

Molecular Weight 61.4 kDa (monomer)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Activity	Measured by its ability to bind MDA-MB-231 human breast adenocarcinoma cells in a flow cytometry assay. A 50-100 fold increase in fluorescence intensity can be observed when MDA-MB-231 cells are stained with 1 µg/mL rhKIR3DL2/Fc Chimera and goat anti-human Fc-PE as compared to cells stained with goat anti-human Fc-PE.
Endotoxin Level	<1.0EU per 1µg of the protein by the LAL method.
SDS-PAGE	85-90kDa, reducing conditions.
Purity	>95%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Reconstitution	100 µg/mL in sterile PBS
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 1 month, 2 to 8°C under sterile conditions after reconstitution. 3 months, -20 to -70°C under sterile conditions after reconstitution.

GENE INFORMATION

Gene Name	KIR3DL2 killer cell immunoglobulin-like receptor, three domains, long cytoplasmic tail, 2 [Homosapiens]
Official Symbol	KIR3DL2
Synonyms	KIR3DL2; killer cell immunoglobulin-like receptor, three domains, long cytoplasmic tail, 2; p140; NKAT4; CD158K; NKAT-4; NKAT4B; killer cell immunoglobulin-like receptor 3DL2; KIR antigen 3DL2; killer Ig receptor; p70 NK receptor CL-5; MHC class I NK cell receptor; CD158 antigen-like family member K; p70 killer cell inhibitory receptor; natural killer-associated transcript 4; natural killer cell inhibitory receptor; p70 natural killer cell receptor clone CL-5; killer cell immunoglobulin-like receptor KIR3DL2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	3812
mRNA Refseq	NM_001242867
Protein Refseq	NP_001229796
MIM	604947
UniProt ID	P43630
Chromosome Location	19q13.4
Pathway	Adaptive ImmuneSystem; Antigen processing and presentation; Graft-versus-host disease; ImmuneSystem; Immunoregulatory interactions between a Lymphoid and a non-Lymphoidcell; Natural killer cell mediated cytotoxicity
Function	receptor activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA