

## Active Recombinant Human CD117 Protein, His&Avi tagged

Cat. No. KIT-62H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	CD117 dimer protein contains a CD117 extracellular domain (UniProt# P10721) fused with a proprietary dimer motif followed by a tandem His-Avi tag at the C-terminus.
<b>Species</b>	Human
<b>Source</b>	HEK293T
<b>ProteinLength</b>	25-524 aa
<b>Description</b>	Human cluster of differentiation 117 (CD117), is a member of the type III receptor tyrosine kinase family. CD117 is also known as KIT, C-Kit, mast/stem cell growth factor receptor (SCFR), KIT proto-oncogene receptor tyrosine kinase, and MASTC. CD117 contains an extracellular domain with five immunoglobulin-like loops, a transmembrane domain, a juxtamembrane domain, and an intracellular domain. CD117 is a type I transmembrane protein, expressed on hematopoietic stem cells, mast cells, melanocytes, germ cells, and interstitial cells of Cajal. CD117 exists as a monomer under normal physical conditions. Upon binding to its natural ligand, stem cell factor (SCF), homodimerization occurs between two CD117 monomers; this homodimerization is essential for its activation. However, oncogenic mutations can cause ligand-independent pathological dimerization and constitutive activation. CD117 is frequently overexpressed or dysregulated in cancers, including gastrointestinal stromal tumors, acute myeloid leukemia, melanoma, and small cell lung cancer. CD117 is a promising drug target,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	especially in precision oncology and regenerative medicine. Understanding CD117 dimerization and its activation is crucial for developing targeted therapeutics.
<b>Molecular Mass</b>	132 kDa
<b>Homodimer/Heterodimer</b>	Homodimer
<b>Purity</b>	Greater than 90% dimer form as determined by SDS-PAGE under non-reducing condition
<b>Application</b>	Verified Applications: ELISA for CD117-specific antibody and stem cell factor (SCF) ligand protein binding assays. Suggested Applications: ELISA for CD117-specific antibody and stem cell factor (SCF) ligand protein binding assays. SPR & BLI for CD117-specific antibody and stem cell factor (SCF) protein binding assays. Animal immunization, RUO.
<b>Storage</b>	At -80 centigrade
<b>Storage Buffer</b>	0.22µm filtered PBS, pH 7.4
<b>Shipping</b>	Frozen Dry Ice


## GENE INFORMATION

<b>Gene ID</b>	3815
<b>Gene Name</b>	KIT v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog [Homo sapiens (human)]
<b>Official Symbol</b>	3815

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Synonyms</b>	KIT; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; PBT, piebald trait; mast/stem cell growth factor receptor Kit; C Kit; CD117; SCFR; p145 c-kit; proto-oncogene c-Kit; piebald trait protein; soluble KIT variant 1; tyrosine-protein kinase Kit; proto-oncogene tyrosine-protein kinase Kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene-like protein; PBT; C-Kit;
<b>mRNA Refseq</b>	<a href="#">NM_000222</a>
<b>Official Symbol 2</b>	<a href="#">KIT</a>
<b>Gene ID 2</b>	<a href="#">3815</a>
<b>Gene Name 2</b>	<a href="#">KIT v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog [Homo sapiens (human)]</a>
<b>mRNA Refseq 2</b>	<a href="#">NM_000222</a>
<b>Protein Refseq 2</b>	<a href="#">NP_000213</a>
<b>MIM 2</b>	<a href="#">164920</a>
<b>UniProt ID 2</b>	<a href="#">P10721</a>
<b>Bioactivity-Antibody Binding</b>	 <p>Activity (cpm)</p> <p>Protein (ng)</p> <p>Immobilized human CD117 dimer protein, His-Avi Tag at 2 µg/mL (100 µL/well)</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	<p>can bind anti-human CD117 monoclonal antibody, with half maximal effective concentration (EC50) range of 4.7-18.8 ng/mL (QC tested).</p>
<p><b>Bioactivity-Ligand Binding</b></p>	<div data-bbox="852 535 1133 842" data-label="Figure"> </div> <p>Immobilized human Stem Cell Factor (SCF) at 2 µg/mL (100 µL/well) can bind human CD117 dimer protein, His-Avi Tag, with half maximal effective concentration (EC50) range of 29.1-116.4 ng/mL (QC tested).</p>
<p><b>SDS-PAGE</b></p>	<p>MW: Molecular Weight marker reduced condition NR: CD117 dimer under non-reducing condition</p>