

Active Recombinant Human Killer Cell Lectin-Like Receptor Subfamily K, Member 1, Fc-tagged

Cat. No. KLRK1-5673H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human KLRK1 protein was expressed in Murine myeloma cell line with the Fc part of human IgG1 Pro100-Lys330 (N-terminus) and Phe78-Val216 (C-terminus).
Species	Human
Source	Mammalian Cells
Protein Length	78-216 a.a.
Description	KLRK1 is a type II transmembrane glycoprotein having an extracellular lectin-like domain. This domain lacks the recognizable calcium-binding sites found in true C-type lectins and binds protein rather than carbohydrate ligands. Human KLRK1 is expressed on CD8+ αβ T cells, γδ T cells, NK cells and NKT cells. In mouse systems KLRK1 also occurs on macrophages. Human ligands for KLRK1 include MICA, MICB, and ULBP1, 2, and 3. Expression of KLRK1 ligands occurs in epithelial cells, tumor cells and under conditions of stress or infection.
Form	Lyophilized from a 0.2 μm filtered solution in PBS.
N-terminal Sequence	Met
Molecular Weight	43 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Activity	Measured by its ability to bind its ligand, rhMICA/Fc Chimera in a functional ELISA.
Endotoxin Level	<1.0 EU per 1 µg of the protein by the LAL method.
SDS-PAGE	50-55kDa, reducing conditions.
Purity	>90%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Reconstitution	100µg/mL in sterile PBS
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 1 month, 2 to 8°C under sterile conditions after reconstitution. 3 months, -20 to -70°C under sterile conditions after reconstitution.


GENE INFORMATION

Gene Name	KLRK1 killer cell lectin-like receptor subfamily K, member 1 [Homo sapiens]
Official Symbol	KLRK1
Synonyms	KLRK1; killer cell lectin-like receptor subfamily K, member 1; KLR; CD314; NKG2D; NKG2-D; D12S2489E; FLJ17759; FLJ75772; NKG2-D type II integral membrane protein; NK cell receptor D; NKG2-D-activating NK receptor; Killer cell lectin-like receptor subfamily K member 1; DNA segment on chromosome 12 (unique) 2489 expressed sequence; OTTHUMP00000239062; OTTHUMP00000239064; CD314 antigen
Gene ID	22914
mRNA Refseq	NM_007360

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_031386
MIM	611817
UniProt ID	P26718
Chromosome Location	12p13.2-p12.3
Pathway	Adaptive ImmuneSystem; Immune System; Immunoregulatory interactions between a Lymphoid and anon-Lymphoid cell; Malaria; Natural killer cell mediated cytotoxicity
PDBrendering based on 1hr.	

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA