

## Recombinant Human KRT19 293 Cell Lysate

Cat. No. KRT19-4875HCL Lot. No. (See product label)

### SPECIFICATION

|                            |   |
|----------------------------|---|
| <b>Species</b>             | Human   |
| <b>Source</b>              | HEK293  |
| <b>Description</b>         | Antigen standard for keratin 19 (KRT19) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.   |
| <b>Components</b>          | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| <b>Size</b>                | 0.1 mg  |
| <b>Storage Instruction</b> | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.   |
| <b>Applications</b>        | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.   |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



## GENE INFORMATION

|                            |   |
|----------------------------|---|
| <b>Gene Name</b>           | KRT19 keratin 19 [ Homo sapiens ]   |
| <b>Official Symbol</b>     | KRT19   |
| <b>Synonyms</b>            | KRT19; keratin 19; keratin, type I cytoskeletal 19; 40 kDa keratin intermediate filament; CK19; cytokeratin 19; K1CS; K19; keratin; type I cytoskeletal 19; type I; 40 kd; MGC15366; CK-19; keratin-19; cytokeratin-19; keratin, type I, 40-kd; 40-kDa keratin intermediate filament; |
| <b>Gene ID</b>             | 3880  |
| <b>mRNA Refseq</b>         | NM_002276   |
| <b>Protein Refseq</b>      | NP_002267   |
| <b>MIM</b>                 | 148020  |
| <b>UniProt ID</b>          | P08727  |
| <b>Chromosome Location</b> | 17q21-q23   |
| <b>Pathway</b>             | Signaling mediated by p38-alpha and p38-beta, organism-specific biosystem;  |
| <b>Function</b>            | protein binding; structural constituent of cytoskeleton; structural constituent of muscle;  |

Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA