

## Active Recombinant Human Kynureninase, Carrier Free

Cat. No. KYNU-11H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant human Kynureninase, expressed in Spodoptera frugiperda, Sf 21 (baculovirus) derived.
<b>Species</b>	Human
<b>Source</b>	Sf21 Cells
<b>Description</b>	<p>Kynureninase is a pyridoxal-5-phosphatedependent enzyme that catalyzes the hydrolytic cleavage of the amino acids L-kynurenine and L-3-hydroxykynurenine to give either anthranilic acid or 3-hydroxyanthranilic acid and alanine. The enzyme is a member of the "kynurenine pathway" enzymes, through which the majority of dietary tryptophan is degraded in the liver, and is involved in the de novo biosynthesis of NAD+. Kynurenine pathway genes are expressed in immune system cells such as macrophages and microglia. During inflammatory responses, the kynurenine pathway in these cells produces quinolinic acid (QA) and not NAD+. QA excites neurons via the activation of NMDA (N-methyl-D-aspartate) receptors resulting in neuronal damage. The tissue damaging property has been demonstrated in AID related dementia complex, Alzheimer's, stroke, epilepsy, and Huntington's disease. Because Kynureninase is one of the key enzymes of QA production, its inhibitors may be useful for the treatment of neurological disorders. The recombinant Kynureninase has been shown to possess specificity for 3-hydroxykynurenine over kynurenine.</p>
<b>Form</b>	Supplied as a 0.2 µm filtered solution in MES and NaCl.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Bio-activity</b>	Measured by its ability to oxidize 3hydroxy kynurenine. The specific activity is >75 pmoles/min/μg, as measured under the described conditions. See Activity Assay Prot centigradeol.
<b>Molecular Mass</b>	51 kDa, reducing conditions
<b>Endotoxin</b>	<1.0 eu per 1 μg of the protein by the lal</1.0
<b>Purity</b>	>95%, by SDSPAGE under reducing conditions and visualized by silver stain.
<b>Usage</b>	FOR RESEARCH USE ONLY
<b>Quality Control Test</b>	12 months from date of receipt, 20 to 70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, 20 to 70 centigrade under sterile conditions after reconstitution.
<b>Warning</b>	Avoid repeated freeze-thaw cycles.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">KYNU kynureninase [ Homo sapiens ]</a>
<b>Official Symbol</b>	KYNU
<b>Synonyms</b>	KYNU; kynureninase; kynureninase (L kynurenine hydrolase); L kynurenine hydrolase; kynureninase (L-kynurenine hydrolase);
<b>Gene ID</b>	<a href="#">8942</a>
<b>mRNA Refseq</b>	<a href="#">NM_001032998</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Protein Refseq</b>	NP_001028170
<b>MIM</b>	605197
<b>UniProt ID</b>	Q16719
<b>Chromosome Location</b>	2q22.2
<b>Pathway</b>	Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of amino acids and derivatives, organism-specific biosystem; NAD biosynthesis II (from tryptophan), organism-specific biosystem; Selenium Pathway, organism-specific biosystem; Tryptophan catabolism, organism-specific biosystem; Tryptophan metabolism, organism-specific biosystem;
<b>Function</b>	hydrolase activity; kynureninase activity; kynureninase activity; kynureninase activity; kynureninase activity; protein homodimerization activity; pyridoxal phosphate binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA