

# Acetyl Histone H3-K9 Quantification Kit (Colorimetric)

Cat. No. Kit-0020 Lot. No. (See product label)

## SPECIFICATION

### Product Overview

Acetyl Histone H3-K9 Quantification Kit (Colorimetric) is use for measuring acetylation of histone H3-K9.

### Description

Acetylation of histones including histone H3 has been involved in the regulation of chromatin structure and recruitment of transcription factors to the gene promoters. Histone acetyltransferases (HAT) and histone deacetylases (HDACs) play a critical role in control of histone H3 acetylation at multiple sites. Histone H3 at lysine 9 (H3-K9) is a primary acetylated site of histone H3. Acetylation of histone H3-K9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Acetylation of histone H3-K9 is tightly involved in the cell cycle regulation, cell proliferation and apoptosis. Acetylation of histone H3-K9 is also an active marker and the balance between histone H3-K9 acetylation and methylation is important for the establishment of specific chromatin structures. An imbalance in the equilibrium of histone H3 acetylation including K9 acetylation has been associated with tumorigenesis and cancer progression. Histone H3-K9 acetylation may be increased by inhibition of HDACs and decreased by HAT inhibition. Thus quantitative detection of acetyl histone H3-K9 would provide useful information for better understanding epigenetic regulation of gene activation and for developing HAT or HDAC-targeted drugs. The Acetyl Histone H3-K9 Quantification Kit (Colorimetric) provides a tool for measuring acetylation of histone H3-K9.

### Applications

For specifically measuring histone H3-K9 acetylation using a variety of mammalian cells (human, mouse, etc.) including fresh and frozen tissues, cultured adherent and suspension cells.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Usage</b>	For research use only (RUO)
<b>Storage</b>	Upon receipt, store C3 and Standard control at $-20^{\circ}\text{C}$ . Store all other components at $4^{\circ}\text{C}$ away from light. The components of the kit should be stable for 6 months when stored properly. Note: Check if buffers C1 and C2 contain salt precipitates before using. If so, warm (at room temperature or $37^{\circ}\text{C}$ ) and shake the buffers until the salts are redissolved.
<b>Kit Components</b>	C1 (10X wash buffer) 20 ml C2 (antibody buffer) 12 ml C3 (detection antibody, 1 mg/ml)* 10 $\mu\text{l}$ C4 (detection antibody, 1 mg/ml)* 10 ml C5 (stop solution) 6 ml Standard control (100 $\mu\text{g/ml}$ )* 20 $\mu\text{l}$ 96 well sample strips (with frame) 96 well standard control strips 3* For maximum recovery of the products, centrifuge the original vial after thawing prior to opening the cap.
<b>Features &amp; Benefits</b>	Quick and efficient procedure, which can be finished within 2.5 hours. Innovative colorimetric assay with no need for radioactivity, electrophoresis, and chromatography. Specifically capturing acetyl H3-K9 with the detection limit as low as 2 ng/well and detection range from 20 ng-5 $\mu\text{g/well}$ of histone extracts. The control is conveniently included for quantification of the amount of acetyl H3-K9. Strip microplate format makes the assay flexible: manual or high throughput. Simple, reliable, and consistent assay conditions

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA