

Collagen Glycation assay Kit (Glucose/fructose)

Cat. No. Kit-0239 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

For a long term examination that researches early saccharification from fructose.

Description

The non-enzymatic reaction of reducing carbohydrates with lysine side chains and N-terminal amino groups of macromolecules (proteins, phospholipids, and nucleic acids) is called the Maillard reaction or glycation. This reaction is initiated by the nonenzymatic reaction of reducing sugars with free amino groups on proteins to form Amadori products. The Amadori products undergo a variety of irreversible dehydration and rearrangement reactions, leading to the formation of advanced glycation end products (AGEs). AGEs have adverse effects on the functional properties of proteins. Many AGEs have fluorescent and covalent cross-linking properties. The accumulation of AGEs is thought to play an important role in the pathogenesis of diabetic patients and the aging process. The collagen that forms bone, skin, and blood vessel is also glycated.

Storage

4°C

Kit Components

1. Collagen Acidic Solution: 5 mL; 2. Neutralizing Solution: 5 mL; 3. Glucose solution (200 mM): 10 mL; 4. Fructose solution (200 mM): 10 mL; 5. Sample Dilution Buffer: 20 mL; 6. Aminoguanidine Solution (20mM): Positive control 0.1 mL

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA