

Adenosine Deaminase Activity Assay Kit (Colorimetric)

Cat. No. Kit-1101 **Lot. No.** (See product label)

SPECIFICATION

Description

Adenosine Deaminase (ADA) (E.C. 3.5.4.4.) is an enzyme that catalyzes the conversion of adenosine and 2'-deoxyadenosine to inosine and 2'-deoxyinosine. Adenosine Deaminase is widely distributed in various tissues and cells. There are two isoforms, ADA1 and ADA2. ADA1 is widely expressed in most cells in the body, particularly in lymphocytes and macrophages. It is present in the cytosol, nucleus and found associated with dipeptidyl peptidase-4 on the cell membrane. ADA2 was first found in the spleen but is predominantly found in the plasma and serum. Increased serum ADA levels are found in certain infectious diseases such as tuberculosis and various liver diseases such as acute hepatitis, alcoholic hepatic fibrosis, chronic active hepatitis to name a few. Adenosine Deaminase is also a marker for T-lymphocyte proliferation. In ADA Activity Assay, inosine formed from the breakdown of adenosine is converted to uric acid with ADA Convertor and ADA Developer. The uric acid is measured at OD 293 nm. The kit measures total Activity of Adenosine Deaminase with limit of quantification of 1 mU recombinant Adenosine Deaminase.

Applications

Detection of Adenosine Deaminase activity

Storage

-20°C

Shipping

Gel Pack

Size

100 assays

Kit Components

ADA Assay Buffer (10x); ADA Convertor; ADA Developer; ADA Substrate; ADA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Positive Control; Inosine Standard (10 mM); U.V. transparent plate (96-well)

Detection method Absorbance (OD 293 nm)

Features & Benefits Rapid, and convenient;
The kit measures total Activity of Adenosine Deaminase with limit of quantification 1 mU recombinant Adenosine Deaminase.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA