

# PPAR $\gamma$ Ligand Screening/Characterization Assay Kit

Cat. No. Kit-2114 Lot. No. (See product label)

## SPECIFICATION

**Product Overview**

The PPAR $\gamma$  Ligand Screening Assay Kit provides a single step fluorescence-based assay for screening potential PPAR $\gamma$ -specific ligands. The assay utilizes the ability of potential PPAR $\gamma$ -binding ligands to displace a fluorescent probe, which has a strong affinity for PPAR $\gamma$  Ligand Binding Domain, resulting in loss of fluorescence of the probe. The relative drop in the fluorescence, as a result of competitive binding of PPAR $\gamma$  ligand, can be correlated to the affinity (and hence IC<sub>50</sub>) of the PPAR $\gamma$  candidate ligand. The PPAR $\gamma$  Ligand Screening Assay Kit is easy to use, faster and more convenient as compared to Fluorescence Polarization and TR-FRET-based screening methods. The assay kit can be used to identify and characterize PPAR $\gamma$ -specific ligands for therapeutic applications.

**Size** 100 assays

**Description**

The Peroxisome Proliferator Activated Receptor (PPAR) family of ligand-activated transcription factors consists of three subtypes encoded by separate genes: PPAR $\alpha$ , PPAR $\delta$  and PPAR $\gamma$ . Of these, PPAR $\gamma$  plays an important role in the regulation of fatty acid storage and glucose metabolism. The genes activated by PPAR $\gamma$  stimulate lipid uptake and adipogenesis by fat cells. Many endogenous molecules such as, polyunsaturated fatty acids like arachidonic acid and its metabolites, are known to bind and activate PPAR $\gamma$ . The binding of activating ligands to the ligand binding domain (LBD) of PPAR $\gamma$  promotes its heterodimerization with retinoic acid-like receptor (RXR), which results in the regulated expression of target genes involved in lipid metabolism. Such ligand-based activation of PPAR $\gamma$  may be responsible for

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	inhibiting the growth of cultured human breast, gastric, lung, prostate and other cancer cell lines. In addition, the thiazolidinedione-based anti-diabetic drugs activate PPAR $\gamma$ with greater specificity than PPAR $\alpha$ .
<b>Applications</b>	Rapid, high-throughput screening of drugs and novel ligands. Development of structure-activity relationship (SAR) models to predict PPAR $\gamma$ /ligand interaction liability of novel compounds.
<b>Target Species</b>	Eukaryotes
<b>Storage</b>	Store kit at -20°C, protected from light. Briefly centrifuge small vials at low speed prior to opening. Read the entire protocol before performing the assay. PPAR $\gamma$ Assay Buffer: Bring to room temperature before use. Store at -20°C. Avoid prolonged storage of the PPAR $\gamma$ Assay Buffer at room temperature or 4°C. Human PPAR $\gamma$ : Store at -80°C. Avoid repeated freeze/thaw cycles. Each vial contains enough protein for 50 assays. PPAR $\gamma$ Assay Probe and Ligand Control: Store at -20°C. Bring to room temperature before use.
<b>Kit Components</b>	PPAR $\gamma$ Assay Buffer: 25 ml PPAR $\gamma$ Assay Probe: 10 $\mu$ l PPAR $\gamma$ (Human Recombinant, 500 $\mu$ g): 2 vials PPAR $\gamma$ Ligand Control (100 mM in DMSO): 10 $\mu$ l 384-well Low Volume Black Plate: 1 Plate
<b>Detection method</b>	Fluorescence (Ex/Em 375/460-470 nm)
<b>Compatible Sample Types</b>	Samples containing drugs, inhibitors or ligands (compounds that can interact and affect PPAR $\gamma$ activity)
<b>Features &amp; Benefits</b>	<ul style="list-style-type: none"> <li>• Simple, highly sensitive, high-throughput compatible</li> <li>• Rapid screening of PPAR<math>\gamma</math> ligands</li> <li>• Kit includes a PPAR<math>\gamma</math> ligand control and a stable, recombinant human PPAR<math>\gamma</math></li> </ul>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA