

Recombinant Human LBP cell lysate

Cat. No. LBP-1853HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human LBP / Lipopolysaccharide Binding derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

LBP lipopolysaccharide binding protein [Homo sapiens]

Official Symbol

LBP

Synonyms

LBP; lipopolysaccharide binding protein; lipopolysaccharide-binding protein; BPI fold containing family D; member 2; BPIFD2; LPS-binding protein; BPI fold containing family D, member 2; MGC22233;

Gene ID

3929

mRNA Refseq

NM_004139

Protein Refseq

NP_004130

MIM

151990

UniProt ID

P18428

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	20q11.23
Pathway	IL6-mediated signaling events, organism-specific biosystem; Immune System, organism-specific biosystem; Innate Immune System, organism-specific biosystem; LPS transferred from LBP carrier to CD14, organism-specific biosystem; Salmonella infection, organism-specific biosystem; Salmonella infection, conserved biosystem; Toll Like Receptor 4 (TLR4) Cascade, organism-specific biosystem;
Function	Gram-negative bacterial cell surface binding; Gram-negative bacterial cell surface binding; Gram-positive bacterial cell surface binding; cell surface binding; lipid binding; lipopolysaccharide binding; lipopolysaccharide binding; lipoteichoic acid binding; protein binding; receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA