

Active Recombinant Human LIF protein, His-tagged

Cat. No. LIF-134H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human LIF(23-202aa) fused with His tag at C-terminal was expressed in Insect cells.
Species	Human
Source	Insect Cells
ProteinLength	23-202 a.a.
Description	LIF, as known as leukemia inhibitory factor, is a pleiotropic glycoprotein belonging to the LI6 family of cytokines. This protein is involved in growth promotion and cell differentiation of different types of target cells, influence on bone metabolism, embryogenesis and inflammation. It is produced by the adrenal cortex and likely enhances its production of cortisol and aldosterone. Also, it can function as an autocrine growth factor in some pancreatic cancers, but induced differentiation in the leukemic cell line M1.
Form	Liquid. In Phosphate Buffered Saline (pH 7.4) containing 10% glycerol.
Bio-activity	Measured in a cell proliferation assay using TF-1 human erythroleukemic cell. The ED50 for this effects is less or equal to 0.5 ng/ml.
Molecular Mass	20.8kDa (189aa) 18-40kDa (SDS-PAGE under reducing conditions.)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	ADPSPLPITP VNATCAIRHP CHNNLMNQIR SQAQLNGSA NALFILYYTA QGEFPNNLD KLCGPNVDF PPFHANGTEK AKLVELYRIV VYLGTS LGNI TRDQKILNPS ALSLHSLNA TADILRGLLS NVLCRLCSKY HVGHVDVTYG PDTSGKDV FQ KKKLGCQLLG KYKQIIAVLA QAFHHHHHH
Endotoxin	< 1.0 EU per 1 microgram of protein (determined by LAL method)
Purity	> 90% by SDS - PAGE
Applications	SDS-PAGE
Storage	Can be stored at +4 centigrade short term (1-2 weeks). For long term storage, aliquot and store at -20 centigrade or -70 centigrade. Avoid repeated freezing and thawing cycles.
Concentration	0.25mg/ml (determined by Absorbance at 280nm)

GENE INFORMATION


Gene Name	LIF leukemia inhibitory factor [Homo sapiens]
Official Symbol	LIF
Synonyms	LIF; leukemia inhibitory factor; CDF; cholinergic differentiation factor; DIA; differentiation inhibitory activity; differentiation inducing factor; hepatocyte stimulating factor III; HILDA; human interleukin in DA cells; D factor; melanoma-derived LPL inhibitor; differentiation-inducing factor; hepatocyte-stimulating factor III; differentiation-stimulating factor; MLPLI;
Gene ID	3976

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_002309
Protein Refseq	NP_002300
MIM	159540
UniProt ID	P15018
Chromosome Location	22q12.2
Pathway	Adipogenesis, organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Direct p53 effectors, organism-specific biosystem; Jak-STAT signaling pathway, organism-specific biosystem; Jak-STAT signaling pathway, conserved biosystem; MicroRNAs in cardiomyocyte hypertrophy, organism-specific biosystem;
Function	cytokine activity; growth factor activity; leukemia inhibitory factor receptor binding; leukemia inhibitory factor receptor binding; receptor binding;



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA