

Recombinant Human LIF, His-GST

Cat. No. LIF-472H Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant human LIF protein, fused to His GST-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques. MW = 47.2 kDa (415aa) confirmed by MALDI-TOF.

Species Human

Source E.coli

Description Leukemia inhibitory factor, also known as LIF, is a pleiotropic cytokine that is expressed by a wide variety of cells including activated T lymphocytes, monocytes, mast cells and neuronal cells. It is involved in the induction of hematopoietic differentiation in normal and myeloid leukemia cells, induction of neuronal cell differentiation, regulator of mesenchymal to epithelial conversion during kidney development, and may also have a role in immune tolerance at the maternal-fetal interface.

Sequences Of Amino Acids

MHHHHHMSPLILGYWKIKGLVQPTRLLEYLEEKYEEHLYERDEGDKWRN
 KKFELGLEFPNLPYYIDGDV KLTQSMAIRYIADKHNLG GCPKERAIEIS
 MLEGAVLDIRYGVSR IAYSKDFETLKVDFLSKLPEMLKMFEDRLCHKTYL
 NGDHVTHPDFMLYDALDVVL YMDPMCLDAF PKLVCFFKRI EAIPQIDKYL
 KSSKYIAWPLQGWQATFGGG DHPPKSDLVPRGSHMSPLPTPVNATCAIR
 HPCHNNLMNQIRSQAQLNGSANALFILYYTAQGEPFPNN LDKLCCGPNVT
 DFPPFHANGTEKAKLVELYRIVVYLGTSLG NITRDQKILNPSALSLSKSL
 NATADILRGLLSNVLCRLCSKYHVGHVDVTYGPDTSGKDV FQKKKLG CQL

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	LGKYKQIIAV LAQAF
Purity	> 90% by SDS – PAGE.
Form	Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 1mM DTT, 20% glycerol, 50mM NaCl.
Concentration	1 mg/ml (determined by Bradford assay).
Storage	Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.
GENE INFORMATION	
Gene Name	LIF leukemia inhibitory factor (cholinergic differentiation factor) [Homo sapiens]
Synonyms	LIF; leukemia inhibitory factor (cholinergic differentiation factor);CDF; D-FACTOR; Emfilermin; HILDA; MLPLI; D factor; Differentiation- stimulating factor; Leukemia inhibitory factor precursor; Melanoma-derived LPL inhibitor; Cholinergic differentiation factor
Gene ID	3976
mRNA Refseq	NM_002309
Protein Refseq	NP_002300
MIM	159540
UniProt ID	P15018

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	22q12.2
Pathway	Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway
Function	Cytokine activity, Growth factor activity, Leukemia inhibitory factor receptor binding, Oncostatin-M receptor binding



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA