

# Active Recombinant Human Lymphotoxin Alpha (TNF Superfamily, Member 1)

Cat. No. LTA-165H Lot. No. (See product label)

## SPECIFICATION

**Product Overview** Recombinant Human lymphotoxin Alpha (TNF Superfamily, Member 1) encoding the human Lymphotoxin alpha (LT-a) protein sequence (containing the signal peptide sequence, and the mature human Lymphotoxin alpha sequence) was expressed in modified *human 293 cells*.

**Species** Human

**Source** HEK293

**Description** Lymphotoxin-alpha (LT-a) is a pro-inflammatory cytokine expressed in activated T cells, B cells, natural killer cells, as well as some non-hematopoietic cells. Activation signals include bacterial products, virus infection, T cell receptor activation, crosslinking of surface immunoglobulin (Ig) on B-lymphocytes, and exposure to UV light in epithelial cells. LT-a regulates a variety of biological processes including cell proliferation, apoptosis, coagulation and lipid metabolism.

**Amino Acid Sequence** LPGVGLTPSAAQTARQHPKMHLAHSNLKPA AHLIGDPSKQNSLLWRANTDRAFLQD  
GFSLSNNSLLVPTSGIYFVYSQVVFSGKAYSPKATSSPLYLAHEVQLFSSQYPFHVPL  
LSSQKMVYPGLQEPWLHSMYHGAAFQLTQGDQLSTHTDGIPHLVLS PSTVFFGAFA  
L.

**Molecular Mass** Lymphotoxin-alpha migrates as a broad band between 20 and 25 kDa in SDS-PAGE due to post-translation modifications, in particular glycosylation. This compares with

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	the unmodified Lymphotoxin-alpha that has a predicted molecular mass of 18.7kDa.
<b>pI</b>	Lymphotoxin-alpha separates into a number of isoforms with a pI between 6 and 9 in 2D PAGE due to post-translational modifications, in particular glycosylation. This compares with the unmodified Lymphotoxin-alpha that has a predicted pI of 8.94.
<b>% Carbohydrate</b>	Purified Lymphotoxin-alpha consists of 0-25% carbohydrate.
<b>Glycosylation</b>	Lymphotoxin-alpha contains N-linked and possibly O-linked oligosaccharides.
<b>Purity</b>	>95%, as determined by SDS-PAGE and visualized by Coomassie Brilliant Blue.
<b>Formulation</b>	When reconstituted in 0.5 ml sterile phosphate-buffered saline, the solution will contain 1% human serum albumin (HSA) and 10% trehalose.
<b>Reconstitution</b>	It is recommended that 0.5 ml of sterile phosphate-buffered saline be added to the vial.
<b>Storage</b>	Lyophilized products should be stored at 2 to 8°C. Following reconstitution short-term storage at 4°C is recommended, and longer-term storage of aliquots at -18 to -20°C. Repeated freeze thawing is not recommended.
<b>Activity</b>	The ED50 of Lymphotoxin-alpha is typically 0.05 - 0.07 ng/ml as measured in acytotoxicity assay using the murine WEHI 164 cell line in the presence of actinomycin D.

## GENE INFORMATION

<b>Gene Name</b>	LTA lymphotoxin alpha (TNF superfamily, member 1) [ <a href="#">Homo sapiens</a> ]
<b>Synonyms</b>	LTA; lymphotoxin alpha (TNF superfamily, member 1); LT; TNFB; TNFSF1;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

lymphotoxin alpha; OTTHUMP00000165897; tumor necrosis factor beta; Tumor necrosis factor ligand superfamily member 1; TNF-beta; LT-alpha

**Gene ID**

4049

**mRNA Refseq**

NM\_000595

**Protein Refseq**

NP\_000586

**UniProt ID**

P01374

**Chromosome  
Location**

6p21.3

**MIM**


153440

**Pathway**


Antigen processing and presentation; Cytokine-cytokine receptor interaction; Type I diabetes mellitus

**Function**


cytokine activity; tumor necrosis factor receptor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

PDB rendering based  
on 1tnr.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA