

Native Peanut Lectin, FITC conjugated

Cat. No. Lectin-1719P **Lot. No.** (See product label)

SPECIFICATION

| | |
|-------------------------|--|
| Product Overview | This product is arachis hypogaea lectin which against Gal beta 1-3 GalNAc (FITC). Affinity purified PNA is conjugated with Fluorescein-5- isothiocyanate (FITC-isomer 1) and purified on Gel filtration column |
| Species | Peanut |
| Source | Peanut |
| Description | <p>Peanut lectin (PNA) is an identical tetrameric carbohydrate free protein with MW of 110 kDa. Thomsen-Friedenreich antigen, T-antigen (Galβ1, 3GalNAc) is present on blood group M & N glycoproteins (after removal of sialic acid with neuraminidase), glyconjugates (Mucin type), gangliosides and many glycolipids. T antigen is rarely expressed on normal colonocytes whereas cells of malignant, premalignant cells express this antigen. Peanut lectin has widely been used to detect T antigen in malignant and premalignant cells. Peanut lectin contains one atom of Ca²⁺ and Mg²⁺ per unit.</p> <p>Inhibiting/Eluting sugars: 200 mM Galactose</p> <p>Carbohydrate-Binding Specificity of PNA: Galβ1, 3 GalNAc>>>>>GalNAc>Gal</p> |
| Form | 10 mM bicarbonate, 150 mM NaCl, pH 8.2, 0.1 mM Calcium chloride and 0.05% sodium azide |
| Applications | ICC/IF. Immunofluorescence 1:250-1:1000. The optimum dilution should be determined by |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



the individual lab.

Storage

Store at 2-8centigrade.

Concentration

5 mg/ml (Please refer to the vial label for the specific concentration)

GENE INFORMATION

Synonyms

PNA lectin, PNA, Arachis hypogaea, Peanut Lectin, Peanut glutinin ctin, Arachis hypogaea, Peanut agglutinin, Arachis hypogaea lectin, Peanut agglutinin lectin

UniProt ID

[P02872](#)

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA