

Native Wheat Germ Lectin, Biotin conjugated

Cat. No. Lectin-1727W **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This product is the triticum vulgaris lectin which agansts N-acetyl glucosamine (Biotin).
Species	Wheat Germ
Source	Wheat Germ
Description	<p>Wheat germ agglutinin (WGA) (MW 36 kDa) is a dimeric, carbohydrate-free protein composed of two identical subunits. The amino acid sequence provides a figure of 21,600, based on 171 amino acids per polypeptide chain. The protein dissociates into monomers in acidic media, pK of the reaction 4. The receptor sugar for WGA is N-acetyl glucosamine (GlcNAc), GlcNAcβ14GlcNAc. These structures are present in many serum and membrane glycoproteins, bacterial cell wall peptidoglycans, chitin, cartilage glycosaminogycons and glycolipids. Native WGA also react with glycoproteins with sialic acid residue. This lectin is useful for the purification of insulin receptors, serum proteins and neuronal tracing.</p> <p>nhibiting/Eluting sugars: 500 mM GlcNAc, dimer, trimmer or 100 mM acetic acid.</p> <p>Carbohydrate-Binding Specificity of WGA: Pentaacetylchitopentaose > Tetraacetylchitotetraose > triacetylchitotriose > Diacetylchitobiose > GlcNAc.</p>
Form	10 mM phosphate, 150 mM NaCl, pH 7.6, 0.1 mM Calcium chloride and 0.05% sodium azide
Applications	ELISA, IHC, WB.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Histochemistry 1: 250-1:1000; WB 1:500-1:2,500; ELISA 1:500-1:2,500. Dilute biotinylated lectin in PBS containing 0.1 mM calcium chloride ions. For Histochemistry the tissues are processed same as for Immunohistochemistry, after blocking step biotinylated lectin is applied followed by streptavidin conjugated to enzyme and chromogen. For WB incubate membrane with blocking protein followed by biotinylated lectin, streptavidin enzyme conjugate and chromogen. For ELISA the target proteins are absorbed on ELISA plate at a concentration of 5-10 µg/ml (50-100 µl), followed by blocking with protein solution, Biotinylated lectin, Streptavidin enzyme conjugate, ELISA substrate. Please refer to Histochemistry, WB and ELISA protocol for detail information. The optimum dilution should be determined by the individual lab.

Storage Store at 2-8centigrade.

Concentration 5 mg/ml (Please refer to the vial label for the specific concentration)

GENE INFORMATION

Synonyms Wheat Germ agglutinin; Triticum vulgaris; Wheat germ agglutinin lectin, Wheat germ agglutinin, Triticum vulgaris, GlcNAc lectin, WGA, WGA Lectin, Triticum vulgaris lectin

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA