

Active Native Dolichos Biflorus Lectin Protein, Fluorescein labeled

Cat. No. Lectin-1770D **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This product is the Fluorescein labeled Dolichos Biflorus Agglutinin and has sugar specificity to N-Acetylgalactosamine. The excitation maximum is at 495 nm and the emission maximum is at 515 nm.
Species	Dolichos Biflorus
Source	Dolichos Biflorus
Description	Dolichos biflorus agglutinin is a glycoprotein with a molecular weight of about 111 kDa and consists of 4 subunits of approximately equal size. This lectin has a carbohydrate specificity toward α -linked N-acetylgalactosamine. It has been used to establish secretor status in blood group A individuals by hemagglutination inhibition techniques and for blood typing. This lectin has also been used as a general marker of developing renal collecting ducts.
Form	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM Ca ²⁺
Bio-activity	Inhibiting/Eluting Sugar: 200 mM N-acetylgalactosamine Carbohydrate-Binding Specificity: N-Acetylgalactosamine.
Molecular Mass	111 kDa
Applications	Immunofluorescence, Glycobiology

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Usage The recommended concentration range for use is 5-20 g/ml.

Storage Refrigerate in the dark. If a precipitate forms upon long-term storage at 4 centigrade, warm to 37 centigrade. and centrifuge before use.

Concentration 2 mg/ml active conjugate

GENE INFORMATION

Synonyms Lectin; DBA

UniProt ID [A0A089ZWN7](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA