

Active Native Peanut Lectin Protein, Rhodamine labeled

Cat. No. Lectin-1817P **Lot. No.** (See product label)

SPECIFICATION

Product Overview	This product is the Rhodamine labeled Peanut Agglutinin (PNA) Lectin and has sugar specificity against Galactose. The excitation maximum is at 550 nm and the emission maximum is at 575 nm.
Species	Peanut
Source	Peanut
Description	Peanut agglutinin binds preferentially to the T-antigen, a galactosyl (β -1,3) N-acetylgalactosamine structure present in many glycoconjugates such as M and N blood groups, gangliosides, and many other soluble and membrane-associated glycoproteins and glycolipids. With certain exceptions, the receptor sequence for PNA is normally sialylated which prevents the lectin from binding to its receptor oligosaccharide (see Jacalin). Even sialic acid which is not bound directly to the receptor sugars may inhibit binding. The presence of calcium ions in diluents can enhance the binding of PNA to receptors, possibly by neutralizing the negative charges on sialic acid residues adjacent to the receptor sequence.
Form	10 mM HEPES, 0.15 M NaCl, pH 7.5, 0.08% sodium azide, 0.1 mM Ca ²⁺
Bio-activity	Inhibiting/Eluting Sugar: 200 mM galactose
Molecular Mass	110 kDa
Applications	Immunofluorescence, Glycobiology

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Usage The recommended concentration range for use is 5-20 g/ml.


Storage Refrigerate in the dark.

Concentration 5 mg/ml

GENE INFORMATION

Synonyms Lectin; PNA

UniProt ID [P02872](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA