

Human Lung Tissue Lysate

Cat. No. Lung-30H Lot. No. (See product label)

SPECIFICATION

Species	Human
Form	The lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% β -mercaptoethanol.
Concentration	2.0 mg/ml
Tissue Type	Lung
Preparation method	Lung tissue lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 μ g/ml of aprotinin, 5 μ g/ml of leupeptin). Tissue and cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay.
Safety Testing	This material has been tested by accepted techniques and has been found to be negative for HBsAg, HIV 1/2, and HCV. Additional patient information may be available upon request.
Diagnosis	Normal
Expiration Date	One year from date of receipt

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Size	0.1 mg, 0.5 mg
Storage Instruction	This lysate should be aliquoted in working amounts and may be stored for up to one month at -20°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles. For longer term storage, freeze working aliquots at -70°C.
Applications	This lysate is for use in Western blotting, 10 µg to 20 µg per lane is recommended for mini gel.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA