

Recombinant Mouse MAP2K4 cell lysate

Cat. No. MAP2K4-557MCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Mouse MAP2K4 / JNKK1 / MKK4 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Map2k4 mitogen-activated protein kinase kinase 4 [Mus musculus]
Official Symbol	MAP2K4
Synonyms	MAP2K4; mitogen-activated protein kinase kinase 4; dual specificity mitogen-activated protein kinase kinase 4; MEK 4; JNKK 1; MAPKK 4; JNK kinase 1; MAPK/ERK kinase 4; SAPK/ERK kinase 1; SAPK/Erk/kinase 1; MAP kinase kinase 4; JNK-activating kinase 1; C-JUN N-terminal kinase kinase 1; mitogen activated protein kinase kinase 4, variant 1; MEK4; MKK4; Sek1; JNKK1; Serk1; PRKMK4;
Gene ID	26398
mRNA Refseq	NM_009157
Protein Refseq	NP_033183
Pathway	Activated TLR4 signalling, organism-specific biosystem; Apoptosis, organism-specific

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem; ErbB signaling pathway, organism-specific biosystem; ErbB signaling pathway, conserved biosystem; FAS pathway and Stress induction of HSP regulation, organism-specific biosystem;

Function

ATP binding; JUN kinase kinase activity; MAP kinase kinase activity; kinase activity; mitogen-activated protein kinase kinase kinase binding; nucleotide binding; protein binding; protein kinase activity; protein serine/threonine kinase activity; protein tyrosine kinase activity; transferase activity; transferase activity, transferring phosphorus-containing groups;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA